SOUTH PLAINS EMERGENCY MEDICAL SERVICES

PRE-HOSPITAL TREATMENT PROTOCOLS for ADVANCED EMT (AEMT)





Charles Addington II, D.O. Medical Director South Plains EMS

> March 1, 2022 Effective Date February 28, 2023 Expiration Date

© 2000, South Plains EMS, Inc.

All Rights Reserved. No part of this document may be used or reproduced, in whole or in part, in any manner, except with written permission of South Plains EMS, Inc., P.O. 53597, Lubbock, Texas 79453-3597, (806) 791-2582.

SPEMS Medical Direction Committee Organizational Chart



The following SPEMS organizations are authorized to utilize these Protocols under the SPEMS Medical Director:

- Amherst EMS
- Associated Ambulance Authority
- Bailey County EMS
- City of Littlefield EMS
- Cochran County EMS
- Crosbyton Clinic Hospital EMS
- Dickens County Ambulance Service
- Earth EMS
- Farwell EMS
- Floydada EMS
- Groom EMS
- Hale Center EMS Association
- Kent County EMS
- King County EMS
- Lazbuddie EMS
- Lubbock Aid Ambulance
- Lynn County Hospital District EMS
- Motley County Ambulance Service
- Olton Volunteer Ambulance Association

- Paducah Ambulance Service
- Paducah Volunteer Fire Department
- Panhandle EMS
- Petersburg EMS
- Plains EMS
- Plainview Fire Department
- Post-Garza County EMS
- Ralls Volunteer Ambulance Service
- Seagraves Loop EMS
- Seminole EMS
- Seminole Volunteer EMT Association
- Silverton EMS
- Springlake Volunteer Fire Department First Responder
- Stinnett EMS
- Sudan Fire Department/EMS
- Wellington EMS
- White Deer EMS
- Wolfforth EMS

PREFACE

These Protocols, originally developed by Francis C. Jackson, M.D., have been revised by the Protocol Committee of South Plains EMS, Inc. The patient care procedures are intended to reflect current recommendations from organizations including, but not limited to, the American Heart Association, The National Association of Emergency Medical Technicians, and the Committee on Trauma of the American College of Surgeons. They are not intended to restrict or substitute for the use of informed professional judgment by the Medical Control Physicians.

To the best of our knowledge, drug dosages are consistent with national standards. Neither the Protocol Committee, the Medical Direction Committee, South Plains EMS, Inc., nor the Medical Directors shall be held liable for readers' errors, omissions, or misunderstanding of the text.

DISCLAIMER

The original version of these protocols are located in the SPEMS office, any changes whatsoever are strictly prohibited without the express written permission of the SPEMS's Medical Director.

On occasion a variance, addendum, or other change may be needed to the current SPEMS protocols. In this event the request MUST be facilitated through the SPEMS office. The request will then be submitted to the SPEMS Medical Director for approval.

TABLE OF CONTENTS

SPEMS Medical Direction Committee Organizational Chart	i.
Preface	
Table of Contents	_P-2
General Guidelines	_P-3
Medical Control Authorization	_P-5
EMS Director's Responsibilities	P-7
Procedural Guidelines for Medical Control Authorization	_ P-7
Infection Control	_P-8
Definitions	P-11
Patient's Physician on Scene	P-14
Unknown Physician on Scene	P-15
	P-17
Pre-Hospital Medications & Intravenous Fluids	_P-33
Decision-Making in Cardiopulmonary Resuscitation	P-34
Do Not Resuscitate Orders	P-36
Cardiac Triage/Transport Decision Scheme	P-43
Stroke Triage/Transport Decision Scheme	-P-44
Pediatric Trauma Triage/Transport Decision Scheme	P-45
Pediatric Burns Triage/Transport Decision Scheme	P-46
Pediatric Non-Trauma Triage/Transport Decision Scheme	P-47
Trauma Triage Criteria for EMS Facility Bypass & Transfer	_P-48
Glasgow Coma Score/Revised Trauma Score	P-50
Guidelines for Trauma Team Activation	_P-51
Trauma Triage/Transfer Decision Scheme – Pre-Hospital	_P-53
Criteria for the Consideration of Air Medical Transport for Trauma Patients	P-54
South Plains EMS Equipment List	_P-55

TREATMENT PROTOCOLS

TRAUMATIC EMERGENCIES	
Burns	1
Trauma	2
RESPIRATORY EMERGENCIES	
Foreign Body Airway Obstruction	4
Near Drowning	
Respiratory Distress	6
CARDIOVASCULAR EMERGENCIES	
Cardiac Arrest/Semiautomatic External Defibrillation	7
Cardiac Chest Pain	8
Cardiogenic Shock	
LVAD	10
Post Resuscitation Management	12
OTHER MEDICAL EMERGENCIES:	
Allergic Reaction	13
Cold Exposure (Systemic Hypothermia)	
Decreased Level of Consciousness	15
Fever/Sepsis	16
Hazardous/Toxic Material Exposure	17
Heat Exposure (Heat Stroke)	18
Hypovolemia (Non-Traumatic)	
Neonatal Resuscitation	20
Obstetric Emergencies	
Poisoning / Overdose	22
Seizures	23
Stroke / TIA or Neurological Deficit	24

GENERAL GUIDELINES

The algorithms in this manual consist of <u>treatment protocols</u>, which may be implemented by EMS technicians at the appropriate level of certification, they establish guidelines based on established standards of care to assist physicians in providing on-line medical control.

Only Advanced EMTs certified by the Texas Department of State Health Services (TDSHS), who also have current medical control authorization from the South Plains Emergency Medical Services (SPEMS) Medical Director, may manage patients under these <u>treatment protocols</u>.

Advanced EMTs who do not have current medical control authorization from the SPEMS Medical Director are prohibited from operating under these <u>treatment protocols</u>.

Execution of <u>treatment protocols</u> without medical control authorization will constitute the unauthorized practice of medicine and may result in action being initiated to revoke the offender's EMS certification.

Medical Direction/Medical Control

Per State laws and DSHS policies, only a **physician** licensed in Texas, may provide medical direction/medical control to EMS personnel. This prevents EMS personnel from accepting orders and directions originating from any non-physician practitioners; including Nurse Practitioners (NPs) and Physician Assistants (PAs).

Medical Control and On-Line Medical Direction for scene calls (Non Interfacility Transfers):

If treatment outside of the SPEMS Protocols is indicated, Medical Control/Medical Direction may be obtained from: • Physician at the receiving facility's emergency department

- In the event that orders are needed and the physician at the receiving hospital cannot be contacted, the medical control physician (ED physician) at University Medical Center shall be contacted immediately.
 NOTE: Obtaining on-line medical direction from the ED physician at UMC does NOT prevent EMS from transporting the patient to the original destination or the destination of the patient's choice.
- In the event that orders are received from the receiving facility that the EMS crew feels may injure or harm the patient, the EMS crew should immediately contact the medical control physician (ED physician) at University Medical Center for guidance.
- In the event that the receiving facility's ED is staffed only with a Nurse Practitioner or Physician's Assistant and orders are needed, the medical control physician (ED physician) at University Medical Center shall be contacted immediately. NOTE: Obtaining on-line medical direction from the ED physician at UMC does NOT prevent EMS from transporting the patient to the original destination or the destination of the patient's choice.
- While treatment procedures can be performed without on-line medical control, it is always available when needed. When in doubt, contact on-line medical control.

Medical Control and On-Line Medical Direction for Interfacility Transfers:

For interfacility transfers, EMS personnel should utilize the following for medical control/medical direction

- If required treatment follows within the SPEMS Protocols, EMS personnel may follow the SPEMS Protocols
- If required treatment is outside SPEMS Protocols, Medical Control/Medical direction may be obtained from:
 - Transferring physician (written and signed orders recommended)
 - $\circ\,$ Receiving physician through verbal or written communications
 - o ER physician of the receiving facility
- Orders from a Nurse Practitioner or Physician's Assistant CANNOT be accepted
- While treatment procedures can be performed without on-line medical control, it is always available when needed. When in doubt, contact on-line medical control.

Medical Control For On-Scene Physicians

If a physician is physically on scene, refer to Patient's Physician on Scene or Unknown Physician on Scene section of these protocols (whichever is applicable).

General Guidelines for Medical Control/On-Line Medical Direction

Radio orders may be communicated through a registered nurse if the physician is physically present in the emergency department but is unable to come to the radio. If all efforts to establish voice contact fail, the authorized AEMT should execute the appropriate protocols and transport. Attempts to establish voice contact with the receiving facility or University Medical Center should continue at least every 5 minutes throughout transport. **NOTE: Obtaining on-line medical direction from the ED physician at UMC does NOT prevent EMS from transporting the patient to the original destination or the destination of the patient's choice.**

Use of these <u>treatment protocols</u> when attempts to make voice contact with the physician have failed, must be communicated to the physician at the receiving facility on arrival, or when contact is made, and must be fully documented by the on the run report. All such instances will be followed by a complete case review.

Use of these treatment protocols must be fully documented by Advanced EMTs on the run report.

These <u>treatment protocols</u> are also intended to serve as guidelines to medical control physicians regarding the current standard for pre-hospital emergency care. These protocols will also provide a basis for auditing the quality of pre-hospital care delivered in the South Plains EMS System. These <u>treatment protocols</u> are not intended to restrict or substitute for the use of informed professional judgment by medical control physicians.

This manual is the result of a review and compilation by the South Plains EMS System Protocol Committee of current, nationally accepted standards for pre-hospital care. The Medical Director of the South Plains EMS System has approved these <u>treatment protocols</u> for use.

This document will be reviewed annually by the Protocol Committee and the Medical Director. Appropriate changes will be distributed following the annual review. If no changes are made, a letter to that effect will be circulated.

These <u>treatment protocols</u> will remain in effect for the duration of the provider license issued by the Texas Department of State Health Services, unless revoked by the SPEMS Medical Director.

MEDICAL CONTROL AUTHORIZATION

Pursuant to the provisions of the Medical Practice Act, the Medical Director of South Plains EMS, Inc. and the SPEMS Medical Direction Committee have determined that the following requirements must be met by Advanced EMTs who wish to operate under the <u>treatment protocols</u> in this manual.

Furthermore, these requirements must also be satisfied by any Advanced EMT who wishes to practice in the SPEMS system under the supervision of the SPEMS Medical Director. This requirement, established pursuant to 22 TAC, Sec. 197.3(b)(1), states that the medical director shall approve the level of pre-hospital care which may be rendered locally by each of the EMS personnel employed by and/or volunteering with the EMS under the Medical Director's supervision before the certificant is permitted to provide pre-hospital care to the public, regardless of the level of state certification.

It is the responsibility of the individual's EMS Service Provider to keep records of individual EMS certifications, immunization records, protocol exams, case review participation, skills exams and any specialized training required by the medical director i.e.: IM injections. These records are subject to examination at anytime, by the SPEMS Medical Director or his/her designate.

Each Advanced EMT must:

- 1. Provide documentation of current certification as an Advanced EMT by the Texas Department of State Health Services. (A current National Registry card can NOT be used in the place of a current TDSHS certification)
- 2. Be currently certified in Health Care Provider BLS.
- 3. Demonstrate approval of local hospital and/or the local medical director by providing a letter which:
 - a) States understanding that the Advanced EMT will be using the SPEMS protocols.
 - b) Acknowledges that the Advanced EMT will be using University Medical Center for on-line medical control under specific circumstances, and
 - c) States the circumstances under which this medical control will be used.
- 4. Have passed the protocol examination within the past year with a grade of 90% or higher. All initial testing will be done via the internet. A copy of the official protocol exam results email must be forwarded to all EMS directors in which the AEMT is on the roster. Each AEMT shall be allowed to take the test twice. Failure of the Protocol exam twice will require the AEMT to contact the SPEMS office for additional testing and remediation. The Medical Director may require additional formal training in the protocols at any time.

NOTE: The Protocol exam is open book and each AEMT is encouraged to utilize the Protocols to complete the exam.

5. Participate in case review at least four times per year. Two of which are recommended by the end of the first half of the year and the remaining two by the end of the second half of the year The SPEMS Medical Director, or his designee, shall attend all case reviews. A written record will be kept of all case reviews. This record will consist of at least a summary of the cases presented and recommendations made for changes in procedure.

Two case reviews per year from EMS services located in the South Plains region that does not utilize the SPEMS Peer Review process, may count towards this requirement with prior authorization of the SPEMS staff, provided that each of the following conditions are met:

- The case review MUST be physician led
- The EMS Director of the SPEMS service must authorize such case reviews
- The individual must request approval from SPEMS staff and obtain the appropriate documentation form from SPEMS staff PRIOR to the case review.
- The EMS Director must maintain proper documentation of these "outside" case reviews using the form that will be furnished by SPEMS staff. The completed forms are subject to inspection upon request

• This does NOT change or alter the skills sign-off requirements for any personnel under SPEMS medical direction A maximum of two such case reviews will be allowed per year. EMS personnel must still obtain the remaining two Case Reviews at a SPEMS service that utilize these Protocols and the SPEMS Peer Review process in order to meet the annual requirements

Exemption: AEMT Is enrolled in a State approved EMT-P class may be exempted from a Case Review while enrolled and/or performing rotations during that quarter. This is subject to the approval of the EMS Director and local EMS services may opt to NOT allow such exemptions. This exemption can only apply to two (2) case reviews in a calendar year. All case review exemptions must be documented on the SPEMS Case Review Exemption form, located on the SPEMS website and will be submitted annually by the EMS Director along with the SPEMS Compliance Checklist. This exemption does not apply to the SPEMS required skills proficiencies. Therefore, all skills proficiencies must be verified per the protocols.

Example: An AEMT is enrolled in an EMT-P course that the classroom runs from January through May and all rotations must be completed by August. That student may be exempted from the first 2 case reviews, with the EMS Director's approval and properly documented.

- 6. Use an approved standard run report form/software, and by the 10th each month send a copy of all run reports to the SPEMS Medical Director through the SPEMS office.
- 7. Documentation is a critical component of pre-hospital care. All documentation must be complete, accurate and done in a timely manner.

A patient contact form, list of medications given, and procedures performed will be given to the patient's nurse prior to leaving the receiving facility.

The full patient care report (PCR) must contain information that is accurate and must completely describe the patient's condition; treatment provided, response to treatment, and any other information that is pertinent to patient care. The full PCR must be completed within one business day of the delivery of the patient and submitted to the receiving facility per rule §157.11.

All SPEMS providers must use an approved standard run report form/software, with copies of all reports submitted, manually or electronically to the SPEMS Medical Director, through the SPEMS office, by the 10th of each month.

- 8. Advanced EMTs must demonstrate proficiency of the following advanced skills at least twice per year between the period of January 1st through June 30th and July 1st through December 31st by:
 - Performing 6 successful IVs, 3 successful endotracheal intubations, 3 King airways, 3 properly performed adult EZ IO placements, 3 properly performed pediatric EZ IO placements, 2 AED, 2 pleural decompressions, and 3 Intramuscular Epinephrine injections (if currently carried by your service) during actual patient care as documented by an observer* approved by the Medical Director;

OR

Participating in at least one formal laboratory evaluation of IV, endotracheal intubation, King airway, properly
performed adult EZ IO placements, properly performed pediatric EZ IO, AED, pleural decompression and
Intramuscular Epinephrine injection (if currently carried by your service) approved by the Medical Director or
an approved designee*;

 Participating in a combination of these options which adequately demonstrates maintenance of proficiency in IV, endotracheal intubation, King airway, properly performed adult EZ IO placements, properly performed pediatric EZ IO, AED, pleural decompression and Intramuscular Epinephrine injection (if currently carried by your service) to the satisfaction of the Medical Director.

(*Skills may be checked off by a SPEMS Peer Reviewer, Associate Medical Director, SPEMS Medical Director, or by a person approved by the SPEMS Medical Director. To be approved, a person must submit to the SPEMS Medical Director a list of qualifications and must receive an endorsement from the Peer Reviewer that performs peer review for that service. All skills will be verified according to the standards listed on the SPEMS website.)

- Participating once per year in a formal laboratory evaluation of pleural decompression conducted by a SPEMS Peer Reviewer, Associate Medical Director, SPEMS Medical Director, or by a person approved by the SPEMS Medical Director
- 9. These protocols shall be utilized under medical direction of the SPEMS Medical Director in the SPEMS/TSA-B area or during routine transfers from one service area to another.

These protocols may also be followed in the performance of **GOOD SAMARITAN** duties when **OFF DUTY** and not responding with any emergency service agency (i.e. EMS, Police, or Fire Dept.). However, in the event that you are **OFF DUTY** and assisting an EMS service that is not under SPEMS medical direction, their online medical direction must be contacted for approval prior to performing any advanced procedures.

10. The intent of these protocols are for EMS professionals to treat patients as they would a member of their own family.

EMS DIRECTOR'S RESPONSIBILITIES

Each EMS Director has the responsibility to assure that each AEMT on their service meets all of the SPEMS Protocol requirements. Specifically, each EMS Director is required to:

- To assure all personnel meet all SPEMS requirements
- To complete and submit the SPEMS Compliance Check List to the SPEMS office by January 31 for the previous calendar year. This check list is available on the SPEMS website
- To maintain records of successful completion of all SPEMS Protocol requirements including skills proficiencies, case reviews, training, testing, etc.
- To remove from active duty any AEMT that fails to meet the requirements as set forth by the SPEMS Protocols.
- To submit copies of any documentation of SPEMS requirements if requested by the SPEMS office or by the Medical Director

PROCEDURAL GUIDELINES FOR MEDICAL CONTROL AUTHORIZATION

Documentation of continuing education, skills proficiency, and case review attendance will be maintained by the EMS technician's training officer. Failure to maintain appropriate documentation may result in the EMS technician's medical control authorization being suspended.

Suspension means that the technician may not practice until completion of all medical control authorization requirements is documented. Suspension may only be removed by the Medical Director, immediately following documentation of all requirements. The fact that an EMS technician's medical control has been suspended does not relieve the technician of the responsibility for meeting all requirements for the subsequent reporting period.

The individual's EMS service director will determine the administrative consequences of suspension of medical control authorization.

Use of the treatment protocols, or practice as an Advanced EMT without current medical control authorization will constitute the unauthorized practice of medicine and may result in action being initiated to revoke the offender's EMS certification.

Upon suspension of medical control authorization, the concerned individual will be notified by a personal letter from the SPEMS Medical Director.

INFECTION CONTROL

GENERAL:

- Each EMS organization participating in SPEMS will designate an individual to act as its Infection Control Officer. The Infection Control Officer will be responsible to the administrative director of the EMS organization and to the Medical Director of SPEMS for ensuring compliance with these procedures.
- 2. Each EMS organization participating in SPEMS should demonstrate compliance with the OSHA Blood Borne Pathogen Rule, "29 CFR, Part 1910,0130," as fully as possible. All EMS personnel should receive formal initial training on the Blood Borne Pathogen Rule. All personnel should complete refresher training annually.
- 3. EMS personnel are strongly encouraged to document immunity to the following diseases by immunization or, when applicable, by history of prior infection:
 - Rubella (German Measles)
 - Red Measles
 - Mumps
 - Hepatitis B
 - Tetanus-Diphtheria
 - Influenza (yearly)
- 4. EMS personnel should be tested annually for tuberculosis unless contraindicated. Positive reactors should be referred to the public health authorities for appropriate follow-up.
- 5. In the unpredictable and uncontrollable pre-hospital environment, it is safest to follow body substance isolation practices, which consider all body substances to be potentially infectious (i.e. "If it's wet, it's bad!"). The following should be considered as potentially infectious:
 - Amniotic fluid
 - Blood
 - Body fluids with visible blood
 - Cerebrospinal fluid (CSF)
 - Feces
 - Nasal secretions

- Pericardial fluidPeritoneal fluid
- Semen
- Semen
- Sputum
- Sweat
- Synovial Fluid

- Tears
- Teeth
- Tissues
- Urine
- Vaginal secretions
- Vomitus
- 6. The routine utilization of exposure control procedures and appropriate Personal Protective Equipment (PPE) by the individual EMS employee/volunteer is essential to the safety of all involved personnel. Its use can help ensure protection from infectious materials to the EMS employee/volunteer, that individual's family members, other members of the EMS department, subsequent patients and the general public.
- 7. The selection and utilization of appropriate Personal Protective Equipment (PPE) should be based upon its ability to provide an impervious barrier between any potentially contaminating body fluids and the EMS employee/volunteer. Each participating SPEMS department is responsible for the supply, repair, cleaning, replacement, and safe disposal of all exposure control-related Personal Protective Equipment. All required PPE should be supplied to that department's personnel and subsequently maintained by the individual department at no expense to the employee/volunteer.

ROUTINE VEHICLE CLEANING:

- 1. All exposed surfaces in the patient compartment will be kept clean with a 1:10 solution of household bleach in water, or a NIOSH-approved hospital germicide that also has tuberculocidal properties. Gloves will be worn during cleaning.
- 2. All reusable hard equipment, spine boards, cervical immobilization devices and cervical collars will be cleaned after each use with hot soapy water, rinsed, disinfected with a germicidal agent and dried. Spine boards and other wooden equipment with exposed splinters should be discarded or sanded and resealed. Gloves will be worn during cleaning.

3. Stock items and medications will be checked at shift change (or as otherwise specified by local policy & procedure) for expiration dates. Materials with the shortest time to expiration should be used first. Expired materials will not be used and will be removed from the vehicle and disposed of properly.

Proper disposal of medications should be made when (1) the container is cracked, (2) the contents are obviously contaminated, (3) the medication has not been stored in accordance with the directions on the label or package insert, or (4) the date has expired.

- 4. Disposable equipment will be used whenever possible. Used disposable equipment that has been contaminated with body fluids will be placed in a sealed and appropriately labeled "Biohazard" container until it can be incinerated.
- 5. Following each use, non-disposable equipment will be washed with hot soapy water, rinsed, disinfected with a tuberculocidal/germicidal agent, and dried. Gloves will be worn during cleaning. If non-disposable equipment cannot be cleaned immediately, it should be placed in a sealed and appropriately labeled "Biohazard" container until it can be properly cleaned.
- 6. After patient contact, priority will be given to spills of blood or other body fluids. All contaminated areas should be cleaned with an appropriate germicidal agent. Gloves will be worn during cleaning.
- 7. After patient contact, stretcher linens will be changed. Used linens will be placed in an impermeable bag or will be double-bagged until they can be removed from the ambulance. Used linens will be removed from the ambulance at the earliest possible time for laundering. Gloves will be worn when handling linens obviously contaminated with body fluids. Bags containing contaminated linens should be labeled "Biohazard."
- 8. Sharp objects will be immediately placed in a puncture-resistant container. Needles will not be recapped, cut, bent, or removed from the syringe. The entire needle-syringe unit should be discarded. When filled, the container will be discarded in accordance with the local medical facility's "Biohazard Waste" policies.

PATIENT CARE PRECAUTIONS:

- 1. Gloves should be worn on every ambulance call and should be applied before patient contact is made. However, the driver of the emergency vehicle should put on his gloves either before he departs for the emergency or immediately upon arrival. He should not attempt to put on gloves while driving. Latex gloves do not provide puncture protection. At an accident scene, leather gloves should be worn over latex gloves.
- 2. Eye protection should be worn when there is a risk of splattering with body fluids. Eyeglasses with plain glass lenses may be used if industrial safety glasses or face shields are unavailable or impractical, but safety glasses/shields with side panels are preferred due to their added protection.
- 3. Mouth-to-mouth breathing should NOT be performed. The pocket mask with one-way valve or a bag-valve-mask should be used for ventilating patients.
- 4. Clothing soiled with blood or body fluids should be changed as soon as practical. It is recommended that a change of clothing, a jump suit or a surgical scrub suit be available on the ambulance for each crewmember. If the crewmember's skin has been contaminated, he should be allowed to remove contaminated clothing and, if needed, shower as quickly as possible.
- 5. Patients should wear a mask if a pathogenic organism could be present in their respiratory secretions. If the patient will not tolerate the mask, or must receive continuous respiratory care precluding a mask, the ambulance crew should wear a mask. Also, the ambulance exhaust fan should be utilized and, weather permitting, the windows opened to increase the exchange of air out of the vehicle. High-risk conditions indicating the wearing of masks are known cases of mumps; measles; chicken pox; active tuberculosis; or meningitis; or fever accompanied by rash, stiff neck, or productive cough.

- 6. Known AIDS patients should wear a mask to protect them from infection. If the patient will not tolerate the mask, or must receive continuous respiratory care precluding a mask, the ambulance crew should wear a mask. They should notify the patient that this is being done to protect the patient from possible infectious organisms.
- 7. Pregnant EMS technicians should avoid providing direct care to known AIDS patients, since many of the patients excrete cytomegalovirus. CMV is known to cause birth defects.
- 8. EMS technicians with known or suspected infectious diseases should avoid providing direct care until a physician determines that there is no risk of transmitting infection to immunocompromised patients.

HAND WASHING/HAND CARE:

- 1. Vigorous scrubbing of the hands with a germicidal soap under running water for 30 seconds will remove or kill most pathogens. Hands should be washed at the beginning and on completion of duty and immediately following each call as soon as gloves are removed. Wearing gloves does not eliminate the need to wash hands.
- 2. Lotion should be applied following hand washing to avoid chapping of the skin, but some lotions can affect the integrity of latex gloves.
- 3. Cuts or other open lesions on the hands or other exposed skin should be covered with a fluid resistant bandage. Bandaging open lesions does not eliminate the need to glove.

EXPOSURE PROCEDURES:

- 1. With routine utilization of appropriate precautions, the risk of needle stick injuries can be significantly reduced. However, in the event that a needle stick does occur, the site should be encouraged to bleed. The site should be cleaned immediately with an alcohol foam and the hands washed thoroughly as soon as possible.
- 2. All cases of possible disease exposure, including a needle stick, should be reported immediately to the personnel at the receiving hospital and to the appropriate supervisor with the EMS technician's organization. The incident should be thoroughly documented on the EMS Agency's or receiving hospital's applicable incident report form.
- 3. The infection control practitioner at each hospital will follow up all cases of exposure of EMS technicians and will advise on appropriate procedures. State law requires this notification.

DEFINITIONS

1. ACUTE CORONARY SYNDROME:

The clinical presentation associated with ACS may be placed into 1 of 3 general categories.

- Classic angina: which refers to the traditional presentation of chest pain (dull, crushing, substernal etc.).
- Atypical presentation: Chest pain which falls short of the typical features is called atypical chest pain. Examples include pain that is sharp, intermittent, in the teeth, neck, shoulder, arm or abdomen etc. Atypical chest pain mostly encompasses females, diabetics, and the elderly.
- Anginal equivalents: consider in higher risk patients: dyspnea, palpations, syncope or near syncope, generalized weakness with no history of a GI bleed or recent fever, and DKA. Anginal equivalents are symptoms not usually associated with classic angina, but are common "atypical symptoms". They are called "equivalents" because they often are the only symptoms the patient manifests during an ischemic cardiac event. An example would be a diabetic with only vomiting and no chest discomfort.

Risk factors include: smoking, hypertension, age, family history of CAD, obesity, stress, and sedentary life style.

The key to forming an accurate impression of chest pain remains in the clinical history. In order to make this impression, one must look at the patient's physical presentation, listen to their story, and be able to compile and interpret all collected information. If the patient's story/presentation, risk factors, 12 lead and vitals sign point to ACS, then EMS personnel should consider the patient a candidate for the Chest Pain/Possible MI Protocol until proven otherwise.

2. ADEQUATE PERFUSION:

- Patient alert and oriented
- Skin warm and dry
- Palpable radial pulses
- Capillary refill < 2 seconds

3. ADULT REFUSAL OF TRANSPORT

An adult, capable of **Informed consent**, that is competent and medically capable, may refuse treatment and/or transport. In such a case, the patient <u>MUST</u> be informed of the potential risk (including death) of such refusal and must sign a patient refusal form. If the patient refuses to sign, you should document the refusal to sign on the paperwork and have a witness sign the refusal. Witnesses in order of preference, may include Police Department, Family Member, Bystander or EMS Crew.

Adults who are **<u>NOT</u>** capable of Informed Consent <u>**MUST**</u> be treated and transported. Local Law Enforcement and/or Medical Control may be contacted for assistance.

In all cases, a SPEMS run report must be completed.

4. ASEPTIC TECHNIQUE:

Aseptic techniques include practices performed just before, during, and/or after any procedure. It is used to reduce the risk of post-procedure infections and to minimize the exposure of the health care providers to potentially infectious microorganisms. The use of aseptic technique is for all procedures, particularly invasive procedures that may break the skin or mucous membrane. This technique includes, but not limited to:

- Antiseptic hand hygiene and proper use of PPE
- Use of appropriate antiseptics to cleanse the area of the patient's body that is in jeopardy of infection/contamination
- Avoid contamination of equipment and medication
- Skin should not be touched after skin antisepsis. If this happens repeat the aseptic technique for that area
- After the insertion of any device through the patients skin the insertion site shall be covered with the appropriate dressing to prevent infection

5. APPROPRIATE LEVEL OF CARE:

Following a complete assessment of the patient, the Advanced EMT should treat the patient at the Advanced EMT protocol level unless impractical or unobtainable. In that event, the Advanced EMT may treat the patient at the EMT-Basic protocol level. An example of this might be: A very close (< 3 minutes) proximity to the receiving hospital where initiating ALS treatment would not affect patient care. Another example: **Oral Glucose** could be administered as opposed to IV **Dextrose** 10%, in the event that an IV is unobtainable. The Advanced EMT should document these occurrences and justifications in the report narrative.

All incidences mentioned above must be peer reviewed

- 6. <u>CENTRAL NERVOUS SYSTEM SIGNS (observe for the following):</u>
 - Level of consciousness
 - Ability to speak
 - Reaction to painful stimuli
 - Pupil size/reaction to light
 - Ability to move extremities
 - Seizures/abnormal posturing

7. CONTACT MEDICAL CONTROL:

Receiving physician at destination hospital or central medical control physician. Notify regarding patient's condition using the following format:

Medical Patients:

- Identify hospital being called, ambulance unit number, name of service, and Med channel
- Age and sex of patient
- Chief complaint
- Vital signs/GCS (including pulse, blood pressure, respirations, pulse ox and level of consciousness)
- Treatment
- Transport Signal, Code and ETA

Trauma Patients:

- Identify hospital being called, ambulance unit number, name of service, and Med channel
- Age and sex of patient
- Mechanism of injury
- Major injuries
- Trauma modifiers
- Vital signs/GCS/RTS (including pulse, blood pressure, respirations, pulse ox and level of consciousness)
- Treatment
- Transport Signal, Code and ETA

8. CONTINUE TO TREAT, MONITOR AND TRANSPORT:

Continue treatment and assessment of vital signs during transport to hospital, including initial vital signs and the vital signs at time of patient transfer.

9. <u>CROUP:</u>

Characterized by inspiratory and expiratory stridor and a seal-bark like cough, it is most common amongst children < 3 years of age. Croup is often preceded by an upper respiratory infection. Respiratory distress, tachypnea, and retractions are also commonly associated with Croup. One of the most distinctive characteristics of croup is the abrupt or sudden onset of the symptoms noted above.

10. DUTY STATUS/GEOGRAPHICAL AREA/GOOD SAMARITAN

These protocols shall be utilized under medical direction of the SPEMS Medical Director in the SPEMS/TSA-B area or during routine transfers from one service area to another.

These protocols may also be followed in the performance of **GOOD SAMARITAN** duties when **OFF DUTY** and not responding with any emergency service agency (i.e. EMS, Police, or Fire Dept.). However, in the event that you are **OFF DUTY** and assisting an EMS service that is not under SPEMS medical direction, their online medical direction must be contacted for approval prior to performing any advanced procedures.

11. ERRORS/DEVIATIONS

All medication errors, and other inadvertent deviations from SPEMS's protocols, require a written Incident Report. These must be reviewed by the Peer Reviewer at the service's next case review.

12. HIGH CONCENTRATION OXYGEN:

Oxygen delivered either by simple face mask or non-rebreather mask at 10-15 liters per minute. If using bag-valvemask, supplemental **oxygen** should be delivered at 15 liters per minute. A demand valve may also be used to deliver high concentration oxygen. A nasal cannula should generally be avoided on patients with significant illness or injury because it does not provide high concentrations of oxygen.

13. <u>INADEQUATE RESPIRATIONS:</u> SHALLOW respirations <10, or >35 per minute.

14. LEVEL I OR II TRAUMA CENTER:

Hospitals with formal designation as a Level I or Level II Trauma Center by the Texas Department of State Health Services and The American College of Surgeons.

15. MEDICATION CONCENTRATIONS/STORAGE

From time to time, the medications included in these protocols may be supplied in concentrations or amounts other than those indicated. Regardless of the particular manner in which medications are supplied, equivalent total amounts must be present, and it is the EMS technician's responsibility to be certain that correct dosages are administered to patients.

Unless specified otherwise, generic and trade name products are considered interchangeable.

It is the responsibility of the individual EMS provider to make sure that all the stocked drugs are stored as per manufacturer's specification. Documentation as to how drugs are stored may be requested by DSHS.

16. MEDICATION STORAGE AND ACOUNTABILITY

All EMS services which utilize SPEMS Medical Direction and/or Protocols will maintain and store all pharmaceuticals as per the manufacture's recommendation. Medications should be arranged in an orderly and organized fashion (i.e. drug bag/box or ambulance compartment) to facilitate quick access as well as limit medication errors. Medications should be stored in an area of the ambulance which limits access to unauthorized individuals. Ambulances licensed at the ALS/MICU level that are operating at the BLS level, as allowed by rule, should have all ALS/MICU medications secured by no less than a numbered zip tie locking system to assure accountability. Narcotics, if carried, should be stored within a locked cabinet.

An EMS service that utilizes narcotics/controlled medications must follow all DEA and SPEMS reporting and tracking requirements. All medications stocked/stored by an EMS provider should be accounted for by utilizing the individual EMS provider's inventory control procedures and/or policies.

17. MULTIPLE PARALLEL PROTOCOLS:

It is understandable that patients may present with multiple problems that require simultaneous treatment. An example is the patient involved in a trauma and having respiratory distress. In the past there has been a reluctance to implement more than one protocol during a patient encounter.

Crews may simultaneously employ multiple protocols when appropriate. However, they must always be cognizant cumulative and contradicting medications. All considerations cannot be presented here, and the occurrence of this necessity should be infrequent. Questions or clarifications should be referred to the service director, peer reviewer, or medical director.

18. NON-ACCIDENTAL TRAUMA:

Non Accidental Trauma (NAT) or Abuse should always be considered in patients in which the patient's condition is inconsistent with the mechanism of injury or history provided. This applies to all age groups; but especially with pediatrics. In the pediatric patient this should be suspected if the patient is exhibiting head trauma symptoms (decreased LOC, seizures, posturing, unequal pupils, vomiting without fever or diarrhea, bradycardia, cardiac arrest or CPR reportedly done). The EMS provider should maintain a high level of suspicion and situational awareness in these cases. The receiving facility should be notified as soon as possible. Notification should be via radio if possible and advise you suspect trauma, and at the receiving facility upon patient delivery.

19. NON-EMS LICENSED/CERTIFIED PERSONNEL

On occasion, licensed or certified healthcare providers (nurses, respiratory therapists, physician assistants, etc.) may accompany the EMS crew in the back of an ambulance if it has been determined by the crew or by the transferring or receiving facility that patient care would be enhanced. The healthcare provider must obtain prior medical direction in event that care is to be rendered.

Physicians may provide patient care as described in the protocols under "unknown physician on scene" (P-15) or "patient's physician on scene" (P-14).

Students may engage in patient care while under the direct supervision of an approved preceptor.

20. NO TRANSPORT CODES

Classification of calls in the SPEMS Region resulting in a patient not being transported should be noted on dispatch records as follows and documented in the patient care report:

- 1. N-1: Unfounded call / False call.
- 2. N-2: Duplicate call.
- 3. N-3: Injury noted, but patient refused transport.
- 4. N-4: Patient accepted treatment but refused transport.
- 5. N-5: EMS refused transport.
- 6. N-6: No injuries.
- 7. N-7: Transported by other means; should **only** be used if the patient left **prior** to EMS arrival or is in the custody of PD and is going to jail.
- 8. N-8: Dead On Scene

ALL CALLS REQUIRE A WRITTEN REPORT. ALL N-3, N-4, AND N-5 CALLS REQUIRE THAT THE SERVICE DIRECTOR OR (DESIGNEE) REVIEW THE WRITTEN REPORT.

21. <u>"OPTIONAL" OR "RECOMMENDED":</u>

The word "optional" or "recommended" will occasionally be seen throughout the protocols in relation to a piece of equipment, specific treatment, or protocol. When "optional" follows a piece of equipment or specific treatment/protocol it is at the discretion of the individual service as to whether or not that equipment will be stocked or treatment performed. When "recommended" follows a piece of equipment or specific treatment/protocol it remains optional but is highly recommended by the medical director and will become mandatory with the next protocol year update.

22. PATIENT'S PHYSICIAN ON SCENE

- 1. Confirm that the physician present is, in fact, the patient's personal physician. Inform the physician that medical control policy requires him to produce verification of his identity and training.
- 2. Inquire as to whether the physician is willing to assume responsibility for the patient's treatment and is willing to physically accompany the patient to the hospital.
- 3. Before initiating any invasive treatment, establish radio contact with MEDICAL CONTROL; and, after advising the medical control physician of the usual information, advise that the patient's physician is present; identify the physician; and have the physician and MEDICAL CONTROL discuss appropriate means of treatment of the patient while you monitor the conversation.
- 4. In the event of any conflict between orders given by the medical control physician and orders given by the physician on the scene, EMS technicians shall permit the patient's physician to personally perform procedures or treatments which conflict with the orders of MEDICAL CONTROL. EMS technicians shall neither participate in nor administer any treatment to the patient under these circumstances, except as ordered by MEDICAL CONTROL.
- 5. Thoroughly document all occurrences.

23. REQUESTING ASSISTANCE AND/OR HIGHER LEVEL OF CARE

If a patient's condition is unstable or deteriorates during transport and the patient would reasonably benefit from the presence of additional personnel, the regional EMS Communications Center or local dispatch should be contacted to coordinate the response of the closest available ALS unit or MICU.

If an EMS Provider **OR** First Responder Organization operating under SPEMS Medical Direction has been requested for assistance **AND/OR** higher level of care by another EMS Provider that is **NOT** under SPEMS Medical Direction, then that SPEMS Provider or Organization may utilize the SPEMS protocols while transporting in the requesting "Non-SPEMS" EMS Provider's ambulance, with prior approval in place. If prior approval is not in place, then the SPEMS provider should contact the "Non-SPEMS" Medical Direction for approval. Be mindful to take any equipment or pharmaceuticals that may not be stocked on the transporting unit. If conflict arises, (i.e. lack of appropriate equipment, disagreement amongst EMS staff, etc) then the requesting "Non-SPEMS" EMS Provider's Medical Direction must be contacted.

If an EMS Provider operating under SPEMS Medical Direction requests assistance **AND/OR** higher level of care from another EMS Provider **OR** First Responder Organization that does **NOT** operate under SPEMS Medical Direction, then the requested "Non-SPEMS" Provider or Organization may utilize their protocols and Medical Direction while transporting in the SPEMS EMS Provider's ambulance. If conflict arises, (i.e. lack of appropriate equipment, disagreement amongst EMS staff, etc) then SPEMS Medical Direction must be contacted.

Air transport should be considered whenever the patient is possibly critically ill or injured and transport by air would expedite the patient's arrival at the hospital. When air transport is requested, the request should be made as soon as the EMS technicians on the scene have performed the primary patient assessment.

The purpose of this section of protocol is to ensure that the best and most efficient patient care is provided.

24. TRANSPORTATION GUIDELINES

If transporting a patient to a receiving hospital out of your coverage area, you should do the following:

- 1. Bring a transfer form completed by the sending hospital.
- 2. Know the name of the receiving physician.
- Request assistance from the receiving area's Advanced Life Support service if the patient's condition is unstable or his/her condition deteriorates while en route. EMS providers will develop written mutual aid agreements with neighboring communities to facilitate coordination of backup responses. Copies of these agreements will be sent to the Regional EMS Communications Center through the SPEMS office.
- 4. Contact Receiving Hospital **as soon as possible prior to arrival**, and re-establish contact any time patient's condition changes. Monitor patient's overall condition and vital signs every 5 to 10 minutes.
- 5. Notify Lubbock EMS dispatch on Med Channel 9 if transporting a patient into the City of Lubbock Code 3. Include sending hospital, city of origin, destination hospital, route traveling, patient severity, transport code, and estimated time of arrival.
- 6. During interfacility transfers SPEMS personnel should operate under the orders of the transferring physician, except were State statute or regulation dictate otherwise such as a physician written DNR. An attempt should be made to contact the transferring or receiving physician to dictate treatment if a decrease in the patient's condition occurs or at the onset of new complications which need to be immediately addressed. If contact is not possible, SPEMS personnel are authorized to operate under the existing SPEMS protocols. Personnel may also contact Medical Control at anytime. Include in your verbal as well as written report any changes in patient condition, orders received or treatment provided. Interfacility transfers include but are not limited to: Hospital to Hospital, Hospital to Specialized care centers and Hospitals to extended care facilities.

7. No Diversion Policy:

Note: This policy has been adopted by TSA-B and applies to all facilities in which SPEMS EMS services transport to; even those outside of the TSA-B region.

- a. Hospitals will no longer be able to divert 911 responses due to ED overcrowding
- b. SPEMS EMS services will honor patient requests to facility of their choice or transport patients to the appropriate facility for their specific needs. If EMS is told to divert, they will only follow this order if it is in the patient's best interest (divert a burn patient to UMC, a pediatric patient to a children's hospital, major trauma to a designated trauma center).
- c. Hospitals will only be allowed to divert EMS in the case of an internal disaster (fire, flood, active shooter, etc.), OR
- d. Facilities who do not have all services may divert based on lack of ability to meet the patient's needs (STEMI, acute stroke or major trauma).
- e. This policy does not apply to EMS services providing interhospital transfers.

25. TRANSPORTATION OF MINORS

A minor is anyone under the age of 18 (with exceptions). Minors may not refuse transport. In those instances where the minor is refusing, and there is no parent or guardian present, Medical Control and/or Law Enforcement should be contacted for assistance.

Exceptions:

- 1. Emancipated from parents.
- 2. Pregnant
- 3. Active duty in the armed forces

26. UNKNOWN PHYSICIAN ON SCENE

- 1. Inform the physician that medical control policy requires him to produce verification of his identity and training.
- 2. Inquire as to whether the physician is willing to assume responsibility for the patient's treatment and is willing to physically accompany the patient to the hospital.
- 3. Before initiating any invasive treatment, establish radio contact with MEDICAL CONTROL; and, after advising the medical control physician of the usual information, advise that a physician is present, identify the physician, and have the physician and MEDICAL CONTROL discuss appropriate means of treatment of the patient while you monitor the conversation.
- 4. In the event of any conflict between orders given by the medical control physician and orders given by the physician on the scene, the medical control physician shall prevail.
- 5. Thoroughly document all occurrences.

Advanced EMT

27. VENTILATION:

Artificial breathing for a patient via mouth-to-mask, bag-valve-mask or demand valve, with high flow **oxygen**. An Oropharyngeal or Nasopharyngeal Airway should be used.

28 VITAL SIGNS:

- Blood Pressure (BP)
- Pulse (rate, regularity, quality)
- Respirations (rate, regularity, quality)
- Skin (color, temperature, moisture)
- Pupil Status (equal/unequal size, round, reactive to light)
- Level of Consciousness (alert, responds to voice or pain, unresponsive)
- Pulse Oximeter

TREATMENT PROCEDURES

AIRWAY MANAGEMENT:

Patients who are apneic, or who are unable to maintain their airway due to severely decreased levels of consciousness, should have an oropharyngeal airway inserted.

Patients who will not tolerate an oropharyngeal airway, but who show signs of inability to maintain an airway without assistance, should have a nasopharyngeal airway placed.

Patients requiring intubation should be pre-oxygenated before attempting endotracheal intubation. Attempts should not exceed 30 seconds without resuming ventilations. Intubation attempts will be limited to 2 attempts by the primary Advanced EMT and 1 attempt by the secondary Advanced EMT/paramedic. An ET tube introducer (i.e. Bougie) may be used, if needed, to facilitate endotracheal intubation at any time. In the event that all 3 attempts at endotracheal intubation fail, secondary airway adjuncts shall be utilized. Patients who cannot be intubated with an endotracheal tube should be intubated with the King airway device. An ET tube introducer (i.e. Bougie) **CAN NOT** be passed through a King airway in order to endotracheally intubate.

Proper placement of the endotracheal tube must be confirmed and must be documented by:

- 1. Observing the tube pass through the glottic opening.
- 2. Observing symmetrical chest rise and auscultating equal breath sounds bilaterally in the lung fields.
- 3. Auscultating for absence of sound over the epigastrium during ventilations.
- 4. Use of pulse oximetry and end-tidal carbon dioxide detection or monitoring.

Proper tube placement should be reconfirmed and documented by auscultation of the lung fields and epigastrium, and observation of the end tidal CO₂ detection monitor following any movement of the patient and upon final disposition at the receiving facility. It is recommended that tube placement also be confirmed by a member of the receiving facility (RN, RT, MD, etc...) prior to turning over patient care to that facility. The confirmation as well as the person confirming the placement of the ET tube should be documented in the run report

Patients who cannot be intubated with an endotracheal tube should be intubated with a King airway (Proper training should be documented prior to use).

BLOOD DRAW FOR LABS:

Due to the importance of rapidly diagnosing Acute Coronary Syndrome and Cerebrovascular Accidents, blood draws will be attempted in the pre-hospital setting. It is required that all units with at least ALS capabilities stock blood tubes to include, but not limited to, "blue top" (PT/PTT INR), "red top (Serology) "purple top" (CBC), and "green top" (BMP/CMP). The tubes listed above (or their equivalent) should be filled appropriately and labeled with the patients first and last name as well as the time the sample was collected. Samples can be drawn on any call as deemed necessary; however, samples should be drawn anytime a CVA or an ACS is in question or when a pt presents with altered mentation. If the IV line is considered to be in jeopardy, then an alternative site should be accessed (i.e. butterfly catheter or Vacu-tainer needle). The order in which the blood tubes must be filled is Blue, Red, Green, Purple.

CENTRAL LINES:

If IV access is impossible and the patient has an existing external venous catheter, contact medical control for permission to access these catheter

CO2 CONTINUOUS WAVEFORM CAPNOGRAPHY

Continuous CO_2 waveform capnography (if available) provides multiple benefits for patients and multiple monitoring aids for advanced EMS providers. The current AHA ACLS guidelines recommend use of continuous quantitative CO_2 waveform capnography for all patients with advanced airways. If available, continuous quantitative CO_2 waveform capnography should be used for the following:

- Continuous confirmation of proper intubation tube placement
 - Goal of ETCO₂ levels between 30 and 34mmHg for perfusing patients
 - Monitoring effectiveness of quality of CPR
 - ETCO₂ readings should be maintained at or above 20mmHg at all times
 - ETCO₂ readings of less than 10mmHg has been shown to have NO chance of return of spontaneous circulation (ROSC)
 - o Readings of less than 20mmHg indicates ineffective CPR
 - If ETCO₂ readings are less than 20mmHg: look at rate, hand placement, depth of compressions, ET tube placement, lung sounds, supplemental oxygen supply, etc.
- Patients with dyspnea, head injuries, or any condition in which the AEMT feels would allow better monitoring and treatment of the patient

CHEMICAL SEDATION / RESTRAINT

Chemical Restraint is a last resort for safely calming extremely agitated patients when the potential for harm to self or others exists. Agitation or acute behavioral disorders may manifest differently. <u>Always suspect an organic cause</u> <u>first</u>. Life-threatening organic conditions that may present with behavioral agitation are hypoglycemia, subdural hematoma, intercerebral hemorrhage, meningitis, hypertensive crisis, and drugs (especially atropine and cyclic antidepressants).

<u>BLS</u>

- 1. Request Law Enforcement assistance on any patient who requires chemical restraint.
- 2. Assess blood glucose to rule-out hypoglycemia.
- 3. Request Paramedic Backup

<u>ALS</u>

- 1. Peripheral intravenous access should be accomplished prior to chemical restraint whenever possible. If unable to obtain due to patient agitation, venous access is to be obtained as soon as possible after the chemical restraint has been safely accomplished.
- 2. Fluid therapy 10-20 ml/kg IV for hypotension (SBP<90mm/Hg). Maintain SBP>90mm/Hg.
- 3. Request paramedic backup for possible chemical restraint.

Excited Delirium Syndrome:

Excited delirium is a medical emergency that has a high risk for increased mortality due to its causes or from the injuries or medical emergencies that may develop secondary to it. Excited delirium needs to be addressed in the prehospital setting, and may involve a combination of psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/bizarre behavior, insensitivity to pain, hyperthermia, and increased strength.

The use of physical restraints and/or tasers may increase the severity of the potential life-threatening condition. In the prehospital setting, excited delirium is most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse; particularly with stimulant drugs such as cocaine, crack cocaine, methamphetamines, amphetamines, or similar agents. Subjects using synthetic marijuana and/or bath salts may also exhibit excited delirium syndrome. Alcohol withdrawals, severe CNS illness, or head trauma may also contribute to the condition.

Management of Excited Delirium Syndrome:

Just as in normal chemical sedation, treatment should occur when there is a danger to the patient or the EMS crew. Since treatment of excited delirium syndrome requires drug administration, Paramedic backup should be requested as soon as possible. An IV should be started if one can safely be established.

GASTRIC TUBE INSERTION:

Gastric tube insertion may be performed in circumstances where gastric distension in a comatose patient is interfering with ventilation. Gastric tube insertion of comatose patients must be preceded by endotracheal intubation to protect against possible aspiration.

HYPERGLYCEMIA

If during routine exam, the patient's BGL is found to be > 300mg/dL, an IV of NS is recommended.

- Patients with vague or non-specific complaints; IV of NS, TKO is recommended.
- If the patient presents severely ill and/or is hemodynamically unstable; IV of NS is recommended to be infused at WO rate for a 1-2 liter bolus or until clinic effect is noted. Pediatric fluid bolus; 20cc/kg or until clinical effect is noted

Caution is advised with infusing high volume crystalloids in patients with CHF or chronic renal failure.

INTRANASAL MEDICATION ADMINISTRATION:

This procedure authorizes the SPEMS Advanced EMT to administer intranasal (IN) medications using the MAD (Mucosal Atomizer Device). The rich vasculature of the nasal cavity provides a direct route into the bloodstream for medications that easily cross the mucous membranes. Due to this direct absorption into the bloodstream, rate and extent of absorption is relatively comparable to IV administration. This procedure allows for the substitution of approved medication administration via the intranasal route in place of other routes listed in specific protocols. This procedure may be used when it is determined by the Advanced EMT that a needleless delivery system may be desired for safety reasons because of patient agitation or combativeness, or an IV is unobtainable. The SPEMS Advanced EMT should refer back to the original protocol upon completion of this treatment guideline. Generally, the intravenous route is the preferred method of drug administration and IV therapy should follow as soon as appropriate or practical.

Required education will be provided at the EMS service level with appropriate documentation of education and skill competency on file.

Indications:

 Adult and Pediatric Altered Level of Consciousness with respiratory compromise. Only the full 2mg dose of Narcan should be administered IN with the exception of the pediatric patient. If the "low dose" of Narcan is warranted it must be administered IV.

Factors that affect absorption (which are relative contraindications):

- Epistaxis
- Facial Trauma
- Nasal Congestion, Discharge or any recognized nasal abnormality
- Destruction of the nasal mucosa from past surgeries or cocaine abuse

Procedure:

- Prepare equipment (MAD, 1 or 3cc syringe, medication, etc.)
- Draw medication in usual fashion allowing for syringe dead space
- •Attach MAD to end of syringe
- •Place the patient in supine or recumbent position if possible
- Stabilize the patients head with one hand as needed
- Press the atomizer against the nostril
- Briskly depress the syringe plunger administering 1/2 of the dose in each nostril
- •Administer maximal dose of 1cc per nostril

Continued on Next Page

General Comments:

- Prior to using a nasal medication, inspect the nostril for significant amounts of blood or mucous discharge. Presence of these will limit the medication absorption. Suctioning the nasal passage prior to delivery may be necessary or an alternative delivery method should be explored.
- Always deliver ½ of the medication dose into each nostril. This doubles the available mucosal surface area (over single nostril) for drug absorption and increases rate and amount of absorption.
- Always use the most concentrated form of the drug available-diluted forms are less effective.
- Do not use over ½ to 1cc of medication per nostril (0.2-0.3cc is the idea volume). If a higher volume is required, apply it in two separate doses allowing a few minutes for the former dose to absorb.
- For small volume doses of medication, be aware that most delivery devices have a "dead space" in the applicator tip where some of the medication will remain. Be sure to take that dead space into account when calculating the volume of medication to be administered. Usually, the addition of 0.1cc of medication will account for the dead space.
- Narcan IN failures are due to being in a hurry to see the patient wake up. IN Narcan takes 3-5 minutes to begin
 working. The patients often just improve breathing so do not always expect full arousal. (The goal is breathing). In
 order to prevent Acute Withdrawal Syndrome, patients that are chronically dependant on opiates should receive
 increments of 0.1mg of Narcan via the intravenous route (IN Narcan can only be administered with the 2mg
 dose with exception of the pediatric patient).

Intranasal Dosages:

DECREASED LOC WITH RESPIRATORY COMPROMISE: Strength: Narcan (Naloxone), 1mg/cc

- **Narcan** (Naloxone) Adult 2mg, IN, may repeat once in 5 minutes if no increase in respiratory rate or LOC.
- **Narcan** (Naloxone) Pediatric 0.1mg/kg, IN, to a max of 2mg, may repeat once in 5 minutes if no increase in respiratory rate or LOC.

MAST:

MAST may be used for splinting of bilateral lower extremity fractures, OR for splinting of unstable pelvic fractures with signs and symptoms of hypovolemia.

NONSPECIFIC COMPLAINT:

If a patient's problem or chief complaint is not addressed by a specific algorithm (e.g. malaise or generalized weakness), the Advanced EMT should initiate appropriate Basic Life Support, perform a thorough patient assessment, and communicate the results of the assessment to Medical Control. If appropriate, an IV, NS, TKO, may be established.

If a patient has profound Nausea/Vomiting, consider administering **Zofran** 4mg IVP. Pediatric (2-12 years of age) dose of **Zofran** is 0.1mg/kg to a max of 4mg, IVP. Do not administer **Zofran** to patients < 2 years of age

OXYGEN THERAPY:

C.O.P.D. patients may be oxygenated using a nasal cannula or a mask at flow rates needed to maintain an **oxygen** saturation of 90-92%. Do not attempt to obtain higher **oxygen** saturations because of the possibility of suppressing the hypoxic drive. Monitor level of consciousness carefully.

To avoid hyperoxia, adult patients with cardiac and/or stroke related problems should be given **oxygen** only if their O2 sats are below 94% or if signs of hypoxia are present. If **oxygen** is required, **oxygen** should be administered, using a non-rebreather, simple face mask or nasal cannula to maintain O2 sats between 94 and 99%. These patients should not be allowed to have a sat of 100%.

All significant trauma patients should receive **oxygen** via non-rebreather or simple face mask to maintain the highest possible **oxygen** saturation.

Except as listed above, all other patients, with an O2 sat below 94%, should receive **oxygen** via non-rebreather or simple face mask. Patient's with an O2 sat of 94% or above; requiring **oxygen** per protocol, may have **oxygen** applied using either a non-rebreather mask, a simple face mask or a nasal cannula.

When a bag-valve-mask is used to ventilate a patient, it should be connected to an **oxygen** reservoir and **oxygen** administered at \geq 15 liters per minute.

PAIN MANAGEMENT FOR TRAUMATIC INJURIES:

- Advanced EMTs may utilize **IV Acetaminophen** ONLY for pain management of **traumatic** injuries when the expected transport time is greater than 20 minutes.
- **IV Acetaminophen** CANNOT be given for any other pain or fever management without approval of on-line medical direction
- **IV Acetaminophen** may be carried in 1,000mg vials and mixed in a 100ml bag of NS or D5W or can be carried in a premixed bag of 1,000mg in 100ml.
- Dosages:
 - o Adult dosage: 1,000mg over 15 minutes via IV infusion (6.67ml/per minute)
 - 67 drops per minute using a 10 drop IV set
 - Pediatric dosages:
 - < 1 month old: 12.5mg/kg over 15 minutes via IV infusion</p>
 - > 1 month old: 15mg/kg to a maximum of 1,000mg over 15 minutes via IV infusion
 - IV Acetaminophen may NOT be repeated without online medical direction orders
- IV Acetaminophen should NOT be administered to any patient with liver disease
- All uses of IV Acetaminophen must be reviewed by the medical director or his designee (peer reviewer)

PARAMEDIC BACKUP:

Whenever a patient would reasonably benefit from MICU care (EKG monitoring, cricothyrotomy, drug therapy or defibrillation), the Regional EMS Communications Center or local dispatch should be contacted to coordinate the response of the closest available MICU.

EMS providers will develop written mutual aid agreements with neighboring communities to facilitate coordination of backup responses. Copies of these agreements will be sent to the Regional EMS Communications Center through the SPEMS office.

The medical control physician may direct a paramedic backup response whenever he believes this would be in the best interest of the patient.

PERIPHERAL VASCULAR ACCESS:

INTRAOSSEOUS INFUSION-ADULT:

Intraosseous infusion is a valuable and safe treatment for the serious or critically ill/injured patient where drug and/or fluid therapy are required for life saving measures. Intraosseous infusion should be considered as a temporary measure during emergencies when other vascular sites are not immediately available. Intraosseous cannulization is not intended for prophylactic use. Only an Advanced EMT with documentation of prior training can perform an Adult IO. If the patient is 40kg and over, and IV access cannot be obtained after three attempts, or 120 seconds, whichever comes first, an EZ-IO intraosseous cannula should be inserted. The preferred site for the EZ-IO is the anterior surface of the proximal tibia. However, the proximal humerus may be utilized as a secondary site if one or more of the contraindications listed below prevents cannulization into the proximal tibia. The EZ-IO must be used as per manufacturer instructions. The EZ-IO may be secured with an EZ-STABILIZER (Optional) as per manufacturer instructions or by other means such as bulky dressings. Manual hand insertion may be done only if the power driver fails. Simply remove the power driver, grasp the needle set and rotate, pushing the needle into the intraosseous space.

Indications:

- 1. Altered mental status (GCS of 8 or less)
- 2. Hemodynamic instability (Systolic BP of < 90)
- 3. Respiratory compromise (respiratory rate of <10 or >35 with low O2 sats not relieved with O2 therapy.
- 4. Profound hypovolemia with altered LOC
- 5. Patient in need of immediate medications or fluids.
- 6. Resuscitation efforts/CPR
- 7. Contact Medical Control with concerns/questions.

Contraindications:

- 1. Previous orthopedic procedure (previous surgery at or around the insertion site, previous IO within 24 hrs.) or injury, or soft tissue injury to the insertion site.
- 2. Fractures of the bone selected for IO infusion.
- 3. Infection at the insertion site.
- 4. Inability to locate landmarks and/or excessive tissue over the insertion site.

Prior to the infusion of fluids or flushes through the IO catheter in the conscious adult patient you may administer **Lidocaine** 1mg/kg SIVP to a max of 50mg if there are no contraindications (Drug allergies, bradycardia, etc...). This will help alleviate the pain in the IO space created by intramedullary pressure due to fluid infusion. Paramedic back up may be requested at any time, but if there is a sudden change in patient condition Paramedic back up must be requested.

Appropriate training and testing must be documented prior to administration of Lidocaine.

The Medical Director or his designee MUST review ALL cases of Adult Intraosseous Cannulization

INTRAOSSEOUS INFUSION-PEDIATRIC:

Intraosseous infusion is a valuable and safe treatment for the serious or critically ill/injured patient where drug and/or fluid therapy are required for life saving measures. Intraosseous infusion should be considered as a <u>temporary</u> measure during emergencies when other vascular sites are not immediately available. Intraosseous cannulization is not intended for prophylactic use. Only an Advanced EMT with documentation of prior training can perform an IO. If the patient is 3-39kg and IV access cannot be obtained after three attempts, or 120 seconds, whichever comes first, an EZ-IO PD intraosseous cannula should be inserted into the anterior surface of the tibia per manufacturer instructions.

The EZ-IO PD may be secured with an EZ-STABILIZER (Optional) as per manufacturer instructions or by other means such as bulky dressings. Manual hand insertion may be done only if the power driver fails. Simply remove the power driver, grasp the needle set and rotate, pushing the needle into the intraosseous space.

Indications:

- 1. Altered mental status (GCS of 8 or less)
- 2. Hemodynamic instability or signs and symptoms of hypoperfusion.
- 3. Respiratory compromise (respiratory rate of <10 or >35 with low O2 sats not relieved with O2 therapy.
- 4. Profound hypovolemia with altered LOC
- 5. Patient in need of immediate medications or fluids.
- 6. Resuscitation efforts/CPR
- 7. Contact Medical Control with concerns/questions.

Contraindications:

- 1. Previous orthopedic procedure (previous surgery of the knee or proximal tibia, previous IO within 24 hrs.) or injury, or soft tissue injury to the insertion site.
- 2. Fractures of the bone selected for IO infusion.
- 3. Infection at the insertion site.
- 4. Inability to locate landmarks and/or excessive tissue over the insertion site.

Prior to the infusion of fluids or flushes through the IO catheter in the conscious pediatric patient you may administer **Lidocaine** 0.5mg/kg SIVP to a max of 50mg if there are no contraindications (Drug allergies, bradycardia, etc...). This will help alleviate the pain in the IO space created by intramedullary pressure due to fluid infusion. Paramedic back up may be requested at any time, but if there is a sudden change in patient condition Paramedic back up must be requested.

Pediatric Bradycardic rates:

- Newborn to 1 yoa < 100bpm
- 1 yoa to 4 yoa < 90bpm
- 5 yoa and older < 60bpm

Appropriate training and testing must be documented prior to the administration of Lidocaine.

The Medical Director or his designee MUST review ALL cases of Pediatric Intraosseous Cannulization

• IV THERAPY:

With the exception of entrapment situations where extrication is required, all IV attempts with unstable trauma patients should take place en route to the hospital. **Two IV attempts** may be performed on non-trauma patients at the scene prior to moving the patient. All other attempts should be made en route to the hospital.

Patients requiring keep open lines may have saline locks placed by personnel trained in this procedure. Precautions should be taken to ensure that the lock and angiocath are flushed with sufficient frequency to maintain patency.

IV fluids should be infused at a wide open rate in all adult cardiac arrest situations to a max of 3000cc.

IV fluids should be infused at 20cc/kg over 10 minutes and may be repeated once in all pediatric arrest situations

PLEURAL DECOMPRESSION:

A 14-gauge X 2 inch or longer IV catheter should be inserted in the mid-clavicular line at the second or third intercostal space. Insert the IV catheter over the superior margin of the rib and withdraw the needle. Stabilize the catheter to avoid kinking. The mid-axillary route should only be used with approval of On-Line Medical Control. When determining the need for a pleural decompression, the AEMT should rely on early signs/symptoms of a tension pneumothorax; primarily absent lung sounds, and should not rely solely on tracheal deviation as tracheal deviation is a late sign.

REMOVAL OF A FOOTBALL HELMET:

In the event an injured football player must be transported, remove the facemask of the helmet to facilitate and secure a patent airway. If it is imperative that the helmet be removed prior to arrival at the emergency department, the shoulder pads SHOULD be removed simultaneously.

RESQPOD (Optional):

The use of the ResQPOD is for the adult patient in cardiac arrest. An adult patient is defined by AHA as one whom has reached puberty. Advanced EMTs who have been properly trained in the use of the ResQPOD may apply the device directly to the ventilation adjunct (i.e. BVM, ET tube, King Airway, etc.). When used with CO₂ monitoring the monitor should be placed between the ResQPOD and the ventilation device. The ResQPOD is not a ventilation device but provides its therapeutic benefit during chest compressions. Therefore, it is necessary to maintain a good seal with the device during the chest compression phase of CPR. In the event that the patient resumes a pulse and/or spontaneous respirations the ResQPOD should be removed.

SEMIAUTOMATIC EXTERNAL DEFIBRILLATION:

Personnel who have received training in the use of the Semiautomatic External Defibrillator may apply and operate the device in cases of cardiac arrest.

SPINAL PRECAUTIONS

Spinal precautions now consist of Spinal Motion Restriction (SMR) rather than full immobilization with a LSB and CIDs.

The use of backboards have not shown any benefit for spinal injuries, and may cause harm. The use of full spinal immobilization is no longer considered the standard of care in most cases.

Therefore, rigid spine boards and similar type devices are generally only used for extrication/transfer devices. Under most instances, the patient should be removed from the LSB or similar rigid device, after transition to the stretcher except when ease of transfer and/or patient safety is a concern; such as morbid obesity. Transporting a patient while remaining on a LSB should be of rare and extraordinary circumstances.

Spinal Motion Restriction consists of the application of an appropriately sized cervical collar and placement on the stretcher and secured to the stretcher with at least three straps. The shoulder straps should be criss-crossed if possible or utilized according to the manufacturer's recommendation. Patients should be transported with head elevated 20 to 30 degrees, unless contraindicated.

Inform the patient to avoid moving his/her head and neck.

Continued on Next Page

If spinal precaution is indicated, per the algorithm or at the attendant's discretion, then Spinal Motion Restriction (SMR) is the preferred technique in MOST cases.

Ambulatory patients who meet criteria for SMR should not be allowed to ambulate further than a few steps and then should be carefully guided to the stretcher and secured. They should not be allowed to step into the ambulance.

Infants and small children: If undamaged and conditions permit, use the car seat with additional padding to reduce spinal motion; when practical.

Multiple Patients: if transporting two or more patients requiring spinal precautions, additional patients can be secured to the bench or captain's seat in the upright position with both the lap and shoulder belt while maintaining spinal motion restrictions. Do not have the patient step up into the ambulance; but utilize the stretcher and then assist the patient to the final position on the bench or captain's seat.

Anytime that a patient that is under Spinal Motion Restriction moves or is moved, manual stabilization of the C-spine should be maintained during movement.

If the patient requires transfer to the transport stretcher via LSB it is highly recommended that the patient be moved from the LSB in a manner that limits spinal manipulation (i.e. log roll, sheet, scoop).

A traditional cervical immobilization device (CID) is not indicated when utilizing Spinal Motion Restriction. However, in the rare instances that a LSB is utilized for full body immobilization and transport a CID is still indicated.

Once a LSB is utilized for the purpose of full body immobilization and/or a cervical collar has been placed, they can only be removed by a physician.

SPINAL PRECAUTION PROCEDURE:

The acronym "NSAIDS" should be used to remember the steps for this procedure.

- N Neuro Exam: Look for any focal deficits such as tingling, reduced strength, or numbness. Verify that there is no language barrier between you and the patient.
 S Significant MOI: Use judgment; Studies have shown that MOI alone should NOT dictate the need for
- Spinal Motion Restriction (SMR).
- A Alertness and Age: Assess that the patient is conscious, alert, and oriented to person, place, time, and event? Confirm the patient is > 5 years of age.
- I *Intoxication*: Assess for any indicators that the person is intoxicated or under the influence (Illicit or Legal Drugs, Alcohol, etc.)
- •D Distracting Injuries: Look for any other injuries that might distract the patient from spinal pain.
- •S *Spinal Exam:* Check for point tenderness along the spinous process of the entire spine (log rolling the patient to assess this is appropriate while manual c-spine is held in order to rule out the need for Spinal Motion Restriction (SMR)).
- Assess for painful range of motion only if the patient had no tenderness to spine (patient should be able to touch their chin to their chest, extend their neck by looking upward, and turn their head from side to side without any midline spinal pain). Discontinue at any time if the patient develops pain or tenderness to ANY area of the spine and immediately perform spinal motion restriction (SMR).

• Once a cervical collar has been placed, it **MUST ONLY BE REMOVED BY A PHYSICIAN**.

If in doubt, **ALWAYS** err on the side of spinal precaution

See flowsheet on following page

SPINAL PRECAUTIONS*



STROKE/TIA (SUSPECTED):

The goal of this protocol is not only to accurately and quickly identify strokes/TIAs in the field, but to identify large vessel occlusions (LVOs) and report findings to receiving facilities immediately. In the event a stroke is suspected, the Cincinnati Stroke Scale (CSS) should be performed. A positive CSS has a 72% probability of being a stroke and of that percentage, approximately 85% are ischemic. All positive CSS findings should be reported to the receiving hospital from the scene. Limit scene time to \leq 10 minutes (document reason if not possible) and consider rapid transport to the appropriate facility, (P-44). To further increase the chances for the best possible patient outcome the VAN assessment for large vessel occlusions should be performed if applicable. The VAN assessment is used in conjunction with the CSS to help determine if the stroke is associated with an LVO. Prehospital recognition and hospital alert can start the process of early, appropriate resource acquisition.

The key to forming an accurate impression is in the physical assessment findings and the clinical history. The following history and signs and symptoms should increase the suspicion of a stroke.

- History: CVA/TIA, Cardiac/Vascular surgeries, Deep vein thrombosis (DVT), Diabetes, Hypertension, CAD, Atrial-Fibrillation, Blood thinners
- Signs and Symptoms: Altered mentation, Weakness/Paralysis, Visual changes, Sensory loss, Aphasia, Dysarthria, Dysphagia, Syncope, Vertigo/Dizziness, Vomiting, Headache, Seizures, Respiratory pattern changes, Hyper/Hypotension, Trouble walking/Unsteady gait

Strokes/TIA can be mimicked by several deferential diagnosis, such as, Seizures/Todd's Paralysis, Hypoglycemia, Tumor, Trauma, Bell's Palsy, Intoxication, Dehydration/Electrolyte imbalances, Infection, Meningitis/Encephalitis.

If possible, inquire about the patients baseline neurological state. Chronic neurological deficits don't necessarily warrant rapid transport to a Stroke Center. Always err on the side of caution if questionable.



Indication:

• The Cincinnati Stroke Scale should be performed on all patients suspected of having a Stroke/TIA including but not limited to, patients with altered level of consciousness and/or neurological deficits.

Contraindications:

- The patient cannot perform the procedure due to a decreased level of consciousness or a life-threatening condition
- The patient has a known language barrier or other deficit that would trigger false-abnormal findings.

Procedure:

- The Cincinnati Stroke Scale (CSS) is a system used to diagnose a potential stroke in the pre-hospital setting. It
 tests three signs for abnormal findings which may indicate that the patient is having a stroke. If any one of the
 three tests show abnormal findings, the patient may be having a stroke and should be transported without delay.
- Step 1 Facial Droop: have the patient smile or show their teeth
 - Normal: Both sides move equally
 - Abnormal: One side of face does not move as well as the other (or at all) so it appears to droop.
 - Step 2 Arm drift: Ask the patient to raise both arms to eye level with their eyes closed and palms facing upward for 10 seconds.

- Normal: The patient can hold both arms up without downward drift.
- Abnormal: The patient is unable to hold both arms up or there is suggestion of new-onset paralysis/weakness.

The VAN Stroke Assessment should be performed on all patients that test positive for arm drift via the Cincinnati Stroke Scale

- Step 3 Speech: Have the person say, "You can't teach an old dog new tricks," or some other simple, familiar saying.
 - Normal: The patient uses correct words with no slurring.
 - Abnormal: The patient slurs the words, gets some words wrong, or is unable to speak.



Indication:

• Utilize the VAN Stroke Assessment anytime there is positive arm drift or motor control deficit via the CSS.

Contraindications:

- The patient cannot perform the procedure due to a decreased level of consciousness or a life-threatening condition
- The patient has a known language barrier or other deficit that would trigger false-positive VAN findings.

Procedure:

- The VAN Stroke Assessment should be performed on all patients with positive findings of motor deficit /arm drift via the Cincinnati Stroke Scale.
 - 1. *Visual Disturbance Assessment:* Face the patient and have them look directly at your nose. The provider will be testing the four quadrants of periphery vision by holding up their hand with a selected number of fingers. The patient should call out the number of fingers they see raised in each quadrant. The provider will assess the periphery vison fields in the following order; right upper, right lower, left upper, and left lower. Assess for eye tracking during this procedure. Ask the patient if they are experiencing any double vision or partial/full blindness.
 - If the patient cannot accurately assess the number of raised fingers in any of the quadrants or reports visual abnormalities, they are VAN positive. The provider should notify the receiving hospital, per protocol. If negative, move on to the next assessment step.
 - 2. Aphasia Assessment: The provider asks the patient to recognize and name two random objects, such as a pen, watch, or button. For the second portion of the assessment, ask the patient to repeat a short sentence, such as, "the early bird gets the worm". The final step in the aphasia assessment asks the patient to follow two simple, separate command, such as "close your eyes", "make a fist", or "wave". This assesses the comprehension/understanding area of the brain.
 - If the patient is unable to correctly perform these assessments, they are VAN-positive. The
 provider should notify the receiving hospital, per protocol. If preformed correctly, move on to the
 next assessment step.

- 3. **Neglect Assessment:** Assess the patient's eyes for a forced gaze to either side. Do the eyes track evenly if following a pen? To perform the next step of the neglect assessment, ask the patient to close their eyes. With their eyes closed the provider then runs their hand over one of the patient's forearms and asks the patient which arm was touched, followed by the opposite forearm, and then final both forearms simultaneously.
 - If the patient was unable to feel sensation to one of their arms, they are VAN positive. The provider should notify the receiving hospital, per protocol.
 - If the patient was able to recognize sensation to each forearm, then they are VAN negative.

Summary:

"VAN Positive" indicates there is motor control deficit <u>AND</u> at least one of the VAN criteria have been met. "VAN-positive" indicates there is an increased probability of a large vessel occlusion. The VAN assessment does not take the place of the CSS. A patient can still have a stroke while remaining "VANnegative". You must advise the receiving hospital of CSS and VAN findings.

Key Points:

- One of the most important items the pre-hospital provider can obtain is the "time of onset" or "last seen normal time", which forms the basis for all treatment decisions. Be very precise in gathering data to establish the time of onset and report it as an actual time (example: "1347" NOT "about 45 minutes ago"). The patient may not be able to receive thrombolytics or other appropriate therapy at the facility unless this vital information has been obtained by EMS. For patients who "woke up and noticed the stroke", the time the patient was last awake will be the reported time of onset. If possible, obtain contact information of a family member or the reporting party.
- Some types of tissue plasminogen activators (tPA) may be administered within 4.5 hours from time of last seen normal if the risks/benefits are considered acceptable after a neurology consult at the hospital. Clot retrieving advancements have extended this time even further for certain cases. The time frame for prehospital "Stroke Activation" is 6 hours from the "last seen normal" time.
- Try to limit scene time to 10 minutes or less. Do not delay transport for non-life sustaining interventions. Transport to the appropriate stroke facility as documented in the Stroke Triage Scheme. (P-44)
- During transport and as time permits both the CSS and VAN assessments should be repeated with additional findings/changes reported to the receiving facility as well as documented on the PCR.
- Be alert for airway problems (swallowing difficulty, vomiting/aspiration) that may coexist with other stroke symptoms.
- Hypoglycemia can present as a LOCALIZED neurologic deficit, especially in the elderly. Obtain a blood glucose measurement as soon as possible and preferably before stroke activation.
- To accommodate the intravenous contrast for specialized CT scans, the IV catheter should be no smaller than an 18g unless the patient's vasculature prevents large bore access. Due to the importance of rapidly diagnosing Stroke/TIA, blood draws should be attempted in the pre-hospital setting. The order in which the blood tubes must be filled is Blue, Red, Green, Purple.

Refer to the Stroke/TIA Transport Decision Scheme Page (P-44)

TASER PROBE REMOVAL:

If an individual's EMS department policy grants EMS staff permission to remove taser probes, the EMS individual **MAY** make a single attempt to remove the probes. If the probes appear to be embedded in the bone, in a sensitive area, or it appears that the removal will be difficult, leave in place and treat as an impelled object. To lessen the risk of a needle stick injury some type of gripping device (hemostats or pliers) should be used to facilitate the removal. The site should then be cleaned and bandaged as appropriate.

TRANSPORTATION:

Transportation to the hospital may begin at any time the EMS technician judges it to be appropriate, even though the words "Continue to Treat, Monitor and Transport" do not appear until the end of each algorithm.

With the exception of entrapment situations where extrication is required, time on the scene with unstable trauma patients should not exceed 10 minutes. If time on scene exceeds 10 minutes, the narrative should include justification. Emphasis should be placed on transporting these patients so they receive definitive hospital care within one hour of the time they were injured.

Since Basic Life Support alone is unlikely to result in the successful resuscitation of cardiac arrest unless followed promptly by Advanced Life Support, patients in cardiac arrest should be transported as soon as possible. The Regional EMS Communications Center or local dispatch should be contacted to coordinate a response by the closest available Mobile Intensive Care Unit.

All patients transported by stretcher must be transported in a DSHS authorized vehicle.

Air transport should be considered whenever its use would expedite a critical patient's arrival at the hospital. The Medical Control physician may order transport of a patient to begin at any time during a treatment procedure.

TRANSPORT TO FREESTANDING EMERGENCY CENTERS:

Under certain circumstances, it may be acceptable for EMS services to transport to a freestanding emergency center. The following guidelines shall be utilized when considering transport to such a facility.

Authorized Freestanding Emergency Center Destinations:

The following freestanding emergency centers have agreed to receive ambulance transport of certain patients. No other standalone emergency center may be utilized.

- Star ER
 - o 7007 Indiana Avenue
 - o Lubbock, TX 79413
 - o Phone: 806-701-4141
- Covenant HOPD
 - o 7905 Milwaukee Avenue
 - Lubbock, TX 79424
 - Phone: 806-368-5837
- Covenant HOPD
 - o 10205 Quaker Avenue
 - o Lubbock, TX 79424
 - Phone: 806-368-8606
- ER Now
 - o 5800 S. Coulter St.
 - o Amarillo, TX 79119
 - o Phone: 806-398-7744
- ER Now
 - $\circ~$ 2101 Coulter St.
 - o Amarillo, TX 79106
 - o Phone: 806-350-7744

Indications for acceptable usage of approved freestanding emergency centers:

- An uncomplicated medical condition with a high likelihood of discharge from the emergency center, such as:
 - o Abdominal pain with normal vital signs
 - o Uncomplicated nausea, vomiting, and/or diarrhea
 - $\,\circ\,$ Flu-like symptoms, sore throat, minor respiratory infection, earache, cough, and/or rash
 - Painful urination
- Uncomplicated musculoskeletal injuries up to and including closed fractures or dislocations (as long as the neurovascular system appears to be intact)
- · Lacerations and/or abrasions not requiring surgical interventions
- Patients in which spinal precautions are not indicated
- Trauma activation is not expected

Contraindications for usage of approved freestanding emergency centers:

- Age: less than 2 years old or older than 64
- STEMI or high-risk chest pain. Chest pain or possible Acute Coronary Syndrome (ACS) in patients over 40 years old
- Cardiac or traumatic arrest
- Uncontrolled airway or patients receiving ventilatory support
- Patients with advanced airways including CPAP, King Airway, and Intubation
- Suspected stroke or stroke-like symptoms
- Altered mental status
- Pregnancy over 20 weeks gestation
- Potential for neurological involvement
- Hemorrhage with wound packing or tourniquet usage
- Any of the following:
 - o Significant penetrating trauma
 - Ischemic extremities
 - Angulated long bone or open fracture; open joint dislocation
 - Spinal motion restriction is indicated
- EMS judgement, need for surgery, or suspected inpatient admission

KEY POINTS:

- Only the above listed FECs can be utilized by this protocol. Transport to any other FEC is NOT allowed.
- If the patient does not meet the indications, with NO contraindications, the patient CANNOT be transported to a FEC. Inform the patient that you cannot meet his/her request and ask that he/she select a hospital-based ED.
- If the patient does meet the above criteria, refer to the algorithm on the next page
- A thorough assessment must be performed prior to considering transport to an approved FEC
- · Contact the FEC early on to allow for the possibility of diversion to another facility
- Medicare and Medicaid are NOT accepted by Star ER and ER Now. Covenant HOPD does accept Medicare and Medicaid. When transporting a patient to Star ER or ER Now, the patient should be informed that they do not accept Medicare and Medicaid as to avoid bills in which the patient is solely responsible for.
- All transports to an approved FEC must be thoroughly documented
- All transports to an approved FEC must be reviewed by a peer reviewer

See Flowchart on Next Page

EMS TRANSPORT TO FREESTANDING EMERGENCY CENTERS (FEC)



TREATMENT FOR SHOCK:

- Assure Airway, Breathing, Circulation, and control of Bleeding (ABCs)
- Insert oropharyngeal/nasopharyngeal airway if patient is unconscious and consider intubation
- Administer high flow oxygen
- Maintain patient's body temperature
- Elevate lower extremities
- Give nothing by mouth
- Fluid resuscitation

TREATMENT FOR SNAKEBITES

- Cool, Calm environment
- Supportive Measures
- Extremity at heart level
- Zero degree elevation
- No Ice or constricting bands

UNCONTROLLED HEMORRHAGE

At times, direct pressure may fail to control bleeding and it may be necessary to utilize other means to assist in bleeding control.

QUICKCLOT

- Must be the heat free formulas
- Combat Gauze, is a 3in X 4yrd roll of heat free hemostatic gauze
- ACS+, is a heat free hemostatic sponge
 - 1. Blot excess blood from wound with gauze pad
 - 2. Apply Hemostatic agent as directed on package
 - 3. Apply FIRM direct pressure to wound for 5 minutes. (if bleeding persists, apply direct pressure for an additional 5 minutes
 - 4. Apply pressure dressing
 - 5. Deliver empty Hemostatic agent packaging to accepting physician

Instructions for the use of QuickClot are printed on its package.

<u>COMMERCIAL TOURNIQUET DEVICE</u>

Tourniquet application may be indicated when life threatening extremity hemorrhage can not be controlled by any other means.

It is required that a commercial tourniquet is used per manufactures instructions, as improvised tourniquets may not be as effective and/or may cause additional harm. The use of QuickClot in conjunction with a tourniquet application is recommended.

- The tourniquet should never be covered by clothing or bandages
- Record the date and time the tourniquet was applied
- If transport time is extended (> 30 minutes) and absence of amputation or continued hypotension/shock the tourniquet may be LOOSENED to assess for bleeding. DO NOT remove the tourniquet. If bleeding continues re-tighten the tourniquet to loss of distal pulses, if bleeding remains controlled leave the tourniquet in place but assure it is loosened to prevent venous occlusion.

PRE-HOSPITAL MEDICATIONS AND INTRAVENOUS FLUIDS

INHALED MEDICATIONS Oxygen Duo-Neb: Consists of 0.5mg Ipratropium Bromide and 3mg of Albuterol Sulfate in 3ml (Page 6, 10, 13) Levalbuterol (Xopenex) (optional) 1.25mg/3ml (Page 6, 13) Racemic Epinephrine 2.25% 11.25mg/0.5ml (optional) (Page 6)

ORAL MEDICATIONS Acetaminophen, 500mg tablets (Page 16) Activated Charcoal, 50g (P-33) Aspirin, 325mg tablet (Page 8, 10, 11) Ibuprofen 200mg tablets (Page 16) Oral Glucose 15g tube (P-11) Liquid Children's Motrin 100mg/5ml (Page 16, 23)

SUBLINGUAL MEDICATIONS Nitroglycerin, 0.4 mg tablet or spray (Page 8, 10, 11)

INTRAMUSCULAR MEDICATIONS Epinephrine Auto-Injector (Adult), 0.3mg/unit (Page 13) Epinephrine Auto-Injector (Pediatric), 0.15mg/unit (Page 13) Epinephrine (1:1,000) 1mg/1cc (Appropriate training and testing must be documented prior to administration) (Page 13) Glucagon 1mg/unit (optional) (Appropriate training and testing must be documented prior to administration) (Page 15, 23)

Services under SPEMS medical direction may carry **Epinephrine Auto-Injectors** to accommodate both adult and pediatric patients **AND/OR Epinephrine** (1:1,000) 1mg/1cc. However, **Epinephrine** (1:1,000) can only be carried if all active ECA's, EMT's and Advanced EMTs are appropriately trained on IM injections (and the standing Allergic Reaction Protocol). This training must be documented including location, date, and time. Documentation must be readily accessible upon inspection

INTRANASAL MEDICATIONS Narcan (Naloxone), 2mg/2cc (P-19, P-20) (Page 10, 15)

INTRAVENOUS FLUIDS Lactated Ringers (LR) Normal Saline (NS)

INTRAVENOUS MEDICATIONS Diphenhydramine (Benadryl), 50mg/1ml (Page 13) Epinephrine (1:10,000), 1mg/10cc (Page 7) Naloxone (Narcan), 2mg/2cc (Page 10, 15) Ondansetron (Zofran) 4mg/2cc (P-20) (Page 8, 10)

PREMIXED IV MEDICATIONS IV Acetaminophen 1,000mg/100cc (P-21) (Page 1, 3)

INTRAOSSEOUS MEDICATIONS Lidocaine 100mg/5cc (Appropriate training and testing must be documented prior to administration) (P-21, P-22) (Page 11)

PREMIXED INTRAVENOUS MEDICATIONS Dextrose 10% (D10W) 250cc bag of NS. Services may carry D50W (25g/50cc) and 250cc bags of NS until current supply of D50W is exhausted or expires (Page 15, 23)

ENDOTRACHEAL MEDICATIONS (Preferred drug route is IV/IO)

If an endotracheal tube has been placed and venous access is delayed, **Naloxone (Narcan)** may be administered by the endotracheal route.

Medications should be administered at 2 times the recommended IV dose. Medications should be administered in a total volume of 10 cc, diluted with normal saline as needed, and introduced directly into the endotracheal tube or through a suction catheter passed beyond the tip of the endotracheal tube. The medication should be followed by 10 quick ventilations with the bag-valve-mask to aerosolize the medication. Chest compressions should be withheld during these ventilations.

DRUGS NOT SPECIFICALLY INDICATED IN PROTOCOLS:

Although not indicated for routine pre-hospital use, the following drugs are included in the authorized medication list and must be directly requested for use at the discretion of **On-Line Medical Control**:

<u>Activated Charcoal</u>: (Dose 1g/kg up to 50g) Activated Charcoal may be indicated for ingestion of medications or other substances. Activated Charcoal should NOT be used in cases of acid or alkali ingestion, if the patient is unable to swallow or has a decreased level of consciousness.
DECISION-MAKING IN CARDIOPULMONARY RESUSCITATION

The current standard of care requires that resuscitation be implemented when two conditions are fulfilled:

- 1. There is the possibility that the brain is viable.
- 2. There is no medically or legally legitimate reason to withhold resuscitation.

RESUSCITATION EFFORTS SHOULD BE WITHHELD ONLY IF A PATIENT IS APNEIC AND PULSELESS AND ONE OF THE FOLLOWING SITUATIONS EXISTS:

- 1. The patient is decapitated.
- 2. Rigor mortis is present.
- 3. Dependent lividity is present.
- 4. Evidence of tissue decomposition is present.
- 5. Massive trauma to the head, neck or thorax, clearly incompatible with life, is present.
- 6. In a multiple casualty situation, there are inadequate numbers of trained personnel to initiate resuscitation while providing life-saving care to other patients.
- 7. A written Texas Department of State Health Services Out of Hospital DO NOT RESUSCITATE (OOH DNR) is available for immediate inspection by the EMS technicians. (Refer to the Do Not Resuscitate Section)
- 8. The patient is wearing the state-approved DNR bracelet and/or necklace bearing the official Out-of hospital DNR logo. (Refer to the Do Not Resuscitate Section)
- 9. Out-of-Hospital DNR Order forms executed in another state or devices authorized by another state as describe in the Do Not Resuscitate Section. (Refer to the Do Not Resuscitate Section)
- 10. The patient's attending physician is at the scene of the emergency and orders the EMS personnel to withhold resuscitation efforts.
- 11. If no written TDSH OOH DNR order is available, the decision to withhold resuscitation efforts may be made by the Emergency Department physician if requested by one persons of the following list if available, in the following priority:
 - a) The patient's spouse;
 - b) A majority of the patient's reasonably available adult children;
 - c) The patient's parents; or
 - d) The patient's nearest living relative.
- 12. The Emergency Department Physician at the receiving hospital orders EMS personnel, via radio, not to initiate or to terminate resuscitation efforts.

In cases involving a request by family members to withhold resuscitation efforts, you should also include documentation of the request, and names and relationships of the persons making the request. <u>ALL</u> EMS personnel present should sign documentation.

NOTE: The patient's private physician, upon learning of EMS involvement in the resuscitation efforts of one of his/her patient's, should contact the receiving Emergency Department's physician and relay any orders to withhold resuscitation efforts to the Emergency Department physician, since EMS cannot accept these "Do Not Resuscitate" orders directly from the private physician via telephone.

If, at anytime, EMS personnel question the legitimacy of the request to withhold resuscitation efforts, or if there are any indications of unnatural or suspicious circumstances, resuscitation efforts should be initiated, but limited to BLS, until such time as Medical Control is contacted and the Emergency Department physician directs otherwise.

IF ANY DOUBT WHATSOEVER EXISTS, RESUSCITATE!

In the pre-hospital setting, EMS Technicians shall not delegate the decision to initiate or withhold resuscitation to other individuals under any circumstances.

Once initiated, BLS and ACLS shall continue until one of the following occurs:

- 1. Effective spontaneous circulation and ventilation have been restored.
- 2. A physician at the receiving medical facility pronounces the patient dead.
- 3. On-line Medical Control orders termination of resuscitation efforts.
- 4. Attending physician arrives on scene and orders termination.
- 5. A legitimate TDSHS Out of Hospital DNR is presented to ambulance personnel. (Refer to the Do Not Resuscitate Section)
- 6. A legitimate Out-of-Hospital DNR Order forms executed in another state is presented to ambulance personnel. (Refer to the Do Not Resuscitate Section)

EXCEPT AS DESCRIBED ABOVE, UNDER NO OTHER CIRCUMSTANCES WILL THE DECISION TO TERMINATE RESUSCITATION BE MADE BY A NON-PHYSICIAN!

EMS personnel should remember that some patients might appear to be dead and not responsive to resuscitation efforts while actually being potential candidates for successful resuscitation. Therefore, if any doubt exists concerning the patient's potential resuscitation by any of the EMS personnel present, resuscitation efforts should be initiated immediately. The following types of patients should receive special consideration for resuscitation since cases have been documented in which these, and other patients, have been successfully resuscitated following the apparent "death" of the patient:

- 1. Hypothermia
- 2. Hypoglycemia
- 3. Acute drug overdoses
- 4. Poisonings
- 5. Pediatric patients
- 6. Drowning
- 7. Unwitnessed (by trained medical personnel) cardiac and/or respiratory arrest

DO NOT RESUSCITATE ORDERS

When dealing with Do Not Resuscitate Orders, the following guidelines shall be followed:

ACCEPT ANY ONE OF THE FOLLOWING AS PROOF OF A VALID OOH DNR ORDER:

Texas Out-of-Hospital Do-Not-Resuscitate Order Form (OOH DNR)

Note: There is now a new TDSHS form for DNRs. Either the new or the old form is acceptable. A copy of both forms is on subsequent pages.

The Texas OOH DNR Order form is a single page form with the Texas DNR logo printed at the top in red or black. The original or a photocopy is acceptable. A copy of these forms is on subsequent pages. The form is considered valid if:

- 1. One of the six sections have been filled out and signed appropriately. **Note**: Electronic signatures are acceptable. (Only one completed section is required)
 - a. Section A: Declaration of the adult person
 - b. Section B: Declaration by legal guardian, agent or proxy on behalf of the adult person is incompetent or otherwise incapable of communication
 - c. Section C: Declaration by a qualified relative of the adult person who is incompetent or otherwise incapable of communication
 - d. Section D: Declaration by physician based on directive to physicians by a person now incompetent or otherwise incapable of communication
 - e. Section E: Declaration on behalf of the minor person
 - f. Section F: Directive by two physicians on behalf of the adult, who is incompetent or unable to communicate and without guardian, agent, proxy or relative. (**Note**: if this section is utilized, no signature of witnesses or notary are required)
- 2. All relevant portions have been completed.
- 3. Signatures of two witnesses or notary are present; except when section F is utilized (No witness or notary signature is required for Section F). **Note**: Electronic signatures are acceptable
- 4. There appears to be no reason to question its authenticity.

Texas OOH DNR Order Bracelet

Two types of OOH DNR Order bracelets are valid:

- 1. A plastic, hospital-type bracelet that is white in color and has the DNR logo printed in red, as is on the front of the OOH DNR Order form. No other identifying information is printed on this bracelet.
- 2. A stainless steel bracelet similar to the "Medic Alert" bracelets, containing the same DNR logo as on the front of the OOH DNR Order form, or the words "Texas Do Not Resuscitate OOH".

When either bracelet is found around the patient's wrist, honor it as if it were a valid OOH DNR Order Form. Do not honor a bracelet that is not attached to the patient. Do not remove the bracelet from the patient, even when the patient is deceased.

Texas OOH DNR Order Necklace

The OOH DNR Order necklace is made of a stainless steel chain, 16 - 18 inches in length with a one-inch diameter disk attached. The disk has the same DNR logo as is on the front of the DNR Order form. When found around the patient's neck, honor this necklace as if it were a valid paper OOH DNR Order form. Do not honor a necklace that is not attached to the patient. Do not remove the necklace from the patient, even when the patient is deceased.

Out-of-State DNR Orders

Personnel may accept Out-of-Hospital DNR Order forms executed in another state or devices authorized by another state, if:

- 1. The order appears to be on an official, state-mandated form.
- 2. The order appears complete (all relevant portions of the form filled in) and valid.
- 3. There appears to be no reason to question the authenticity of the DNR Order form or device.

Should there be a question regarding an out-of-state DNR Order, initiate resuscitation and contact an on-line medical control physician.

DOCUMENTS THAT MAY NOT BE ACCEPTED:

Do Not Accept:

- 1. Do Not Resuscitate Orders that do not reasonably appear to be on an official, state-mandated form.
- 2. Advanced Directives, Directives to Physicians, Living Wills, A Physician's DNR Order in any form not noted above (such as one written by a physician, physician's assistant, or a nurse practitioner).

Texas OOH DNR Orders and Out-of-State DNR Orders Should Not be Honored when:

- 1. A competent patient, including a competent minor, communicates to EMS personnel a desire to revoke an OOH DNR Order.
- 2. A person having a Durable Power of Attorney for Health Care for the patient or the attending physician, legal guardian, parent (if a minor), or qualified relative, as defined in the TDSHS OOH DNR form, communicates to EMS personnel a desire to revoke an OOH DNR Order.
- 3. The patient is pregnant.
- 4. The patient cannot be conclusively identified as the patient named on the OOH DNR Order form.
- 5. There is an airway obstruction.
- 6. Unnatural or suspicious circumstances are present; including suicide attempt.

If doubt exists as to whether an OOH DNR Order should be honored, initiate resuscitation until:

- 1. A valid OOH DNR Order is found. NOTE: If a valid OOH DNR is presented, after resuscitation has been initiated, resuscitation efforts may be discontinued so long as the validity is not in question or one of the 5 preceding conditions are not found.
- 2. A Medical Control physician orders that resuscitation be stopped, or
- 3. Patient care is transferred to a higher level.

COMPLIANCE WITH OOH DNR ORDER:

NOTE: OOH DNR applies only <u>AFTER</u> the cessation of spontaneous respirations or circulation or in the judgment of the pre-hospital provider, the moment of death is at hand.

- 1. If the patient is found in or develops cardiac and/or respiratory arrest, honor the OOH DNR Order by withholding CPR, placement of advanced airway devices (including ET tube and King Airway), artificial ventilation, placement of the AED, manual defibrillation, and transcutaneous cardiac pacing.
- 2. If an OOH DNR Order is found or presented after the patient assessment and/or treatment has begun, stop the resuscitative treatment immediately even if a positive response has occurred.

- 3. If an OOH DNR Order appears to be valid and the patient is not in cardiac or respiratory arrest, provide care directed toward providing comfort, such as opening the patient's airway, providing oxygen, IV fluids or medications, or any other treatment needed except for advanced airway placement, artificial ventilation, defibrillation, and cardiac pacing. NOTE: Assisting ventilations for a breathing patient with a BVM device is <u>NOT</u> a violation of an OOH DNR and should be performed if needed.
- 4. The original OOH DNR Order form or a photocopy may be honored.
- 5. If the patient is transported, the OOH DNR Order form must accompany the patient; the bracelet or necklace must be on the patient.
- 6. The original or a photocopy of the form should be kept and filed with the pre-hospital patient care report.

DOCUMENTATION:

When a patient in cardiac or respiratory arrest is encountered and an OOH DNR Order form is presented, the following must be documented on the pre-hospital patient care report:

- 1. An assessment of the patient's condition.
- 2. Whether or not the OOH DNR Form was honored. If the form was not honored, a full explanation of the reasons and circumstances must be documented
- 3. The type OOH DNR Order (form, bracelet or necklace) used to confirm the DNR status.
- 4. Any problems regarding implementing the DNR Order, including on scene revocation.
- 5. The name of the patient's attending physician from the OOH DNR form.
- 6. The original or a photocopy of the form should be kept and filed with the pre-hospital patient care report.

SUMMARY

- ONLY the TDSHS OOH DNR or other state (other than Texas) issued DNR may be accepted
- OOH DNR applies to out-of-hospital settings including ERs, Nursing Homes, Physician's offices, clinics, dialysis centers, private residences, etc.
- OOH DNR applies only AFTER the cessation of spontaneous respirations or circulation

EXCEPT: Airway obstruction

Suspicious Circumstances

Suicide, homicide, or other unnatural causes of death

Pregnant patients

Patient or guardian state desire not to follow DNR

- Interventions to be withheld are:
 - CPR
 - Advanced Airways (Intubation and King Airway)
 - Artificial ventilation (does not pertain to assisting ventilations on a breathing patient)
 - Defibrillation (includes AED)
 - Transcutaneous cardiac pacing
- If uncertain, err on side of resuscitation until status can be clarified
- The OOH DNR device (form, bracelet, or necklace) should be left attached to and transported with the patient
- Out-of-state DNR may be honored if no reason to question the authenticity of the order or device exists

STO	RESUSCITATE		tive immediately on the date of executi		
Ders	ons who sign the form m		thority or the document is revoked. Co	omfort measures will be given as no	eded.
Pris	ono ono orga the form h	use sign again under n		Mole/Female (× 1.0.1
	Patient's full legal name — printee	or typed	Date of Birth:	Male/Female (Circle One)
CO	MPLETE ONE OF THE FO	UR BOXES: A, B, C, or D	D. If using Box A, B, or C, Witnes	sses and Physician's Statement	must be completed.
A.	CPR, including the treatm	ents listed below, and I di	ult capable of making an informed rect that none of the following res utaneous Cardiac Pacing, Defi	uscitation measures be initiate	d or continued:
1	Signature		Date	Printed or Typed Name	
B.	otherwise unable to mail am the patient's: 🗌 leg	ake his or her wishes kr al guardian; 🗌 agent unde	r Medical Power of Attorney; 🗌	or Qualified Relative (see back	
	I am acting under the I am acting upon the k I am acting in the pati I am acting in the pati I direct that none of th	guidance of a prior Directi nown values and desires o ent's best interest based up e following resuscitatio		ent's physician. Itinued on behalf of the pat	
	-				
C.	with a terminal or irre I am the minor patient's: I direct that none of th	versible condition. Parent;] legal guardia e following resuscitatio	Date ed by a person acting on beha m; or managing conservator. n measures be initiated or cor	tinued on behalf of the pat	ient: Cardiopulmo-
C.	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th	versible condition. Parent;] legal guardia e following resuscitatio	ed by a person acting on beha m; or 🗌 managing conservator.	lf of a minor patient who h itinued on behalf of the pat	ient: Cardiopulmo-
	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature	versible condition. Parent; legal guardia e following resuscitatio R), Transcutaneous Carc	ed by a person acting on beha n; or managing conservator. n measures be initiated or cor liac Pacing, Defibrillation, Adv	lf of a minor patient who h itinued on behalf of the pat anced Airway Management	ient: Cardiopulmo-
	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CPI Signature SSES: (see qualifications on	versible condition. Parent: legal guardia e following resuscitatio R), Transcutaneous Carc reverse) We have witness	ed by a person acting on beha m; or managing conservator. n measures be initiated or cor liac Pacing, Defibrillation, Adv 	lf of a minor patient who h ttinued on behalf of the pat anced Airway Management Printed or Typed Name	ient: Cardiopulmo-
	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature SSES: (see qualifications on Witness 1 Signature	versible condition. Parent; legal guardia e following resuscitatio R), Transcutaneous Carc reverse) We have witness Date	ed by a person acting on beha n; or managing conservator. n measures be initiated or cor fliac Pacing, Defibrillation, Adv Date ed all of the above signatures. Witness Printed o	If of a minor patient who h ntinued on behalf of the pat ranced Airway Management Printed or Typed Name	ient: Cardiopulmo-
TNES	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CPI Signature SSES: (see qualifications on Witness 1 Signature Witness 2 Signature IAN'S STATEMENT: 1, th	versible condition.	ed by a person acting on beha m; or managing conservator. n measures be initiated or cor liac Pacing, Defibrillation, Adv 	If of a minor patient who h ntinued on behalf of the pat anced Airway Management. Printed or Typed Name r Typed Name r Typed Name med above. I have noted the ex	ient: Cardiopulmo- Artificial Ventilatio
TNES	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CPI Signature SSES: (see qualifications on Witness 1 Signature Witness 2 Signature IAN'S STATEMENT: 1, th	versible condition.	ed by a person acting on beha m; or managing conservator. n measures be initiated or cor diac Pacing, Defibrillation, Adv 	If of a minor patient who h ntinued on behalf of the pat anced Airway Management. Printed or Typed Name r Typed Name r Typed Name med above. I have noted the ex	ient: Cardiopulmo- Artificial Ventilatio
TNES	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature SSES: (see qualifications on Witness 1 Signature Witness 2 Signature IAN'S STATEMENT: I, th nt's medical records, and I Date Only use this box if the otherwise unable to ma I attest to issuance of The patient's specific these circumstances of I direct that none of th	versible condition. Parent; legal guardia e following resuscitatio R), Transcutaneous Carc reverse) We have witness Date Date e undersigned, am the atte direct out-of-hospital healt Physician's signature order is being complet ke his or her wishes kn an Out-of Hospital DNR ti wishes are unknown, but re r are otherwise not in the e following resuscitatio	ed by a person acting on beha m; or managing conservator. n measures be initiated or cor diac Pacing, Defibrillation, Adv 	If of a minor patient who h etinued on behalf of the pat ranced Airway Management Printed or Typed Name r Typed Name r Typed Name ned above. I have noted the es th this order as presented. In behalf of an adult who is is al guardian, agent, or quali munication; OR: nable medical judgement, cons thinned on behalf of the pat	ient: Cardiopulmo- Artificial Ventilatio
YSIC.	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature SSES: (see qualifications on Witness 1 Signature Witness 2 Signature IAN'S STATEMENT: I, th nt's medical records, and I Date Only use this box if the otherwise unable to ma I attest to issuance of The patient's specific these circumstances of I direct that none of th	versible condition. Parent; legal guardia e following resuscitatio R), Transcutaneous Carc reverse) We have witness Date Date e undersigned, am the atte direct out-of-hospital healt Physician's signatore order is being complet ke his or her wishes kn an Out-of Hospital DNR t wishes are unknown, but re r are otherwise not in the e following resuscitatio	ed by a person acting on beha m, ormanaging conservator. n measures be initiated or cor fliac Pacing, Defibrillation, Adv 	If of a minor patient who h etinued on behalf of the pat ranced Airway Management Printed or Typed Name r Typed Name r Typed Name ned above. I have noted the es th this order as presented. In behalf of an adult who is is al guardian, agent, or quali munication; OR: nable medical judgement, cons thinned on behalf of the pat	ient: Cardiopulmo- Artificial Ventilatio
YSIC.	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature SSES: (see qualifications on Witness 1 Signature IAN'S STATEMENT: I, th nt's medical records, and I Date Only use this box if the otherwise unable to ma I attest to issuance of The patient's specific these circumstances of I direct that none of th nary Resuscitation (CP) Signature	versible condition.	ed by a person acting on beha m; or managing conservator. n measures be initiated or con diac Pacing, Defibrillation, Adv 	If of a minor patient who h ttinued on behalf of the pat ranced Airway Management. Printed or Typed Name rTyped Name rTyped Name med above. I have noted the ex th this order as presented. nted name h behalf of an adult who is i al guardian, agent, or quali nunication; OR: nable medical judgement, cons thinued on behalf of the pat vanced Airway Managemen Printed or Typed Name	ient: Cardiopulmo- Artificial Ventilatio
YSIC. paties D.	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature SSES: (see qualifications on Witness 1 Signature Witness 2 Signature IAN'S STATEMENT: I, th nt's medical records, and I Date Only use this box if the otherwise unable to ma I attest to issuance of The patient's specific these circumstances of I direct that none of th nary Resuscitation (CP Signature Signature	versible condition.	ed by a person acting on beha m; ormanaging conservator. n measures be initiated or cor fliac Pacing, Defibrillation, Adv 	If of a minor patient who h etinued on behalf of the pat ranced Airway Management Printed or Typed Name r Typed Name r Typed Name ned above. I have noted the es th this order as presented. neted name n behalf of an adult who is i al guardian, agent, or quali munication; OR: nable medical judgement, cons printed or Typed Name Printed or Typed Name Printed or Typed Name	ient: Cardiopulmo- Artificial Ventilatio
YSIC. paties D.	Only use this box if the with a terminal or irre I am the minor patient's: I direct that none of th nary Resuscitation (CP) Signature SSES: (see qualifications on Witness 1 Signature Witness 2 Signature IAN'S STATEMENT: I, th nt's medical records, and I Date Only use this box if the otherwise unable to ma I attest to issuance of The patient's specific these circumstances of I direct that none of th nary Resuscitation (CP Signature Signature	versible condition.	ed by a person acting on beha m; or managing conservator. n measures be initiated or con diac Pacing, Defibrillation, Adv 	If of a minor patient who h etinued on behalf of the pat ranced Airway Management Printed or Typed Name r Typed Name r Typed Name ned above. I have noted the es th this order as presented. neted name n behalf of an adult who is i al guardian, agent, or quali munication; OR: nable medical judgement, cons printed or Typed Name Printed or Typed Name Printed or Typed Name	ient: Cardiopulmo- Artificial Ventilatio

The Older Version of the TDSHS Out-of-Hospital (OOH) DNR Form: Front Side

Page 2 of 2

Figure: 25 TAC §157.25 (h)(2)

OUT-OF-HOSPITAL DNR INSTRUCTIONS

PURPOSE:

This form was designed to comply with the requirements as set forth in Chapter 166 of the Health and Safety Code (H&SC) relating to the issuance of Out-of-Hospital Do-Not-Resuscitate (DNR) orders for the purpose of instructing Emergency Medical Personnel and other health care professionals to forgo resuscitation attempts and to permit the patient to have a natural death with peace and dignity. This order does NOT affect the provision of other emergency care including comfort care.

APPLICABILITY:

This form applies to all health care professionals operating in any out-of-hospital setting to include hospital outpatient or emergency departments and physician's offices.

IMPLEMENTATION:

A competent adult may execute or issue an Out-of-Hospital DNR Order. The patient's attending physician will document the existence of the directive in the patient's permanent medical record.

If an adult patient is capable of providing informed consent for the order, he/she will sign and date the out-of-hospital DNR order on the front of this sheet in Box A. In the event that an adult patient is unable to provide informed consent, his/her Legal Guardian, agent under Medical Power of Attorney, or Qualified Relative may execute the order by signing and dating the form in Box B. If an adult patient is unable to provide informed consent and none of the persons listed in Box B are available, the treating physician may execute the order using Box D with the consent of a second physician who is not treating the patient and/or is a member of the health care facility ethics committee or other medical committee.

The following persons may execute an out-of-hospital DNR order on behalf of a minor; the minor's parents, the minor's legal guardian or the minor's managing conservator. A person executing a DNR order on behalf of a minor may execute the order by signing and dating the form in Box C. An out-of-hospital DNR order may not be executed unless the minor has been diagnosed by a physician as suffering from a terminal or irreversible condition.

The form must be signed and dated by two witnesses except when executed by two physicians only (Box D).

The original standard Texas Out-of-Hospital DNR form must be completed and properly executed. Duplicates may be made by the patient, health care provider organization or attending physician as necessary. Copies of this completed document may be used for any purpose that the original may be used and shall be honored by responding health care professionals.

The presence of a Texas DNR identification device on a person is sufficient evidence that the individual has a valid Out-of-Hospital DNR Order. Therefore, either the original standard form, a copy of the completed standard form, or the device is sufficient evidence of the existence of the order.

For information on ordering identification devices or additional forms, contact the Texas Department of State Health Services at (512) 834-6700.

REVOCATION:

The Out-of-Hospital Do-Not-Resuscitate Order may be revoked at ANY time by the patient **OR** the patient's Legal Guardian/Agent/Managing Conservator/Qualified Relative, Parent (if a minor), or physician who executed the order. The revocation may involve the communication of wishes to responding health care professionals, destruction of the form, or removal of all or any Do-Not-Resuscitate identification devices the patient may possess.

AUTOMATIC REVOCATION: This Out-of-Hospital DNR order is automatically revoked if the patient is known to be pregnant or in the case of unnatural or suspicious circumstances.

DEFINITIONS:

Attending Physician: The physician who is selected by or assigned to a patient who has primary responsibility for a person's treatment and care and is licensed by the Texas State Board of Medical Examiners or who is properly credentialed and holds a commission in the uniformed services of the United States and who is serving on active duty in this state (H&SC 166.002 (3) & (12))

Qualified Relatives: Those persons authorized to execute or issue an out-of-hospital DNR order on behalf of a person who is comatose, incompetent, or otherwise mentally or physically incapable of communication under Section 166.088 H&SC Section 166.088 refers to 166.039; "One person, if available, from one of the following categories, in the following priority...: (1) The patient's spouse; (2) the patient's reasonably available adult children; (3) the patient's nearest living relative."

Health Care Professional: Means physicians, nurses, physician assistants and emergency medical services personnel, and, unless the context requires otherwise, includes hospital emergency department personnel. (H&SC 166.081 (5))

Witnesses: Two competent adult witnesses must sign the form acknowledging the signature of the patient or the person(s) acting on the patient's behalf (except when signed by two physicians in Section C). Witness One must meet the qualifications listed below. Witness Two may be any competent adult. Witness One (the "qualified" witness) may not be: (1) person designated to make a treatment decision for the patient; (2) related to the patient by blood or marriage; (3) entitled to any part of the estate; (4) be a person who has a claim against the estate of the patient; (5) the attending physician or an employee of the attending physician; (6) an employee of a health care facility in which the patient is being cared for, if he or she is involved in providing direct patient care to the patient; or (7) an officer, director, patner, or business office employee of a health care facility in which the patient is being cared for or any parent organization of the health care facility.

Please report any problems with this form to the Texas Department of State Health Services at (512) 834-6700.

Revised July 19, 2005 Texas Department of State Health Services

Page 2 of 2 Publications No. EF01-11421

The Older Version of the TDSHS Out-of-Hospital (OOH) DNR Form: Back Side

			STATE HEALTH		the second s
DO NOT RESUSCITATE	This document becomes effective immediately on the person is pronocated dead by authorized rand				unhi
Person's full legal name			Date of birth		Maler Female
	son: I am competent and at least 18 years of CPR), transcutaneous cardiac pacing, defib				tinued for me:
Person's signature	er ny dansed aneous cardiac pacing, densi	inación, actancea an	Date	Printed name	
rel solt s sidinardire					
Declaration by legal guardia	n, agent or proxy on behalf of the adult per				Incompations or ethomation
am the: 🔲 legal guardian;	agent in a Medical Power of Attor	nen inen	fally or physically incapable		
	he person, or a determination of the best inter Itation (CPR), transcutaneous cardiac pacin				tiated or continued for the
lignature		Date	Pi	inted name	
Declaration by a gualified color	tive of the adult person who is incompetent	er offerniles bernah	le of communications (or	the shore material manuales	
spouse. F adult child.	parent, OR r nearest living relative, a			The second s	6.088
e person or a determination of th	is incompetent or otherwise mentally or phys the best interests of the person, I direct that no bus cardiac pacing, defibrillation, advanced	ne of the following re	suscitation measures be in		
gnature	and the second	Date		ted name	
Declaration by physician base erson's attending physician and h	ed on directive to physicians by a person no	w incompetent or non	written communication t	o the physician by a competent p	erson: I am the above-noted
	nave: Visued directive to physicians by the adult, now inco	mpetent: OF	obserVed his/her issuance be	ore two witnesses of an OCH-DNR in a no	nwitten manne.
lirect that none of the following Ivanced airway management, a	g resuscitation measures be initiated or con artificial ventilation	tinued for the person	cardiopulmonary resusci	tation (CPR), transcutaneous card	líac pacing, defibrillation,
ttending physician's	Noncal Vestal Station	Date	Printed		Lic
gnature		Diric	name		¢
Declaration on behalf of the n	ninor person: I am the minor's: 👘 📋 pare	ent: 📋 legal gu	ardian: OR 🛛	managing conservator.	
	inor as suffering from a terminal or irreversible				or continued for the person
	(CPR), transcutaneous cardiac pacing, defib	rillation, advanced air	way managément, artific Date	al ventilation.	
agnature			Date		
Printed name					
	ons on backside.) We have witnessed the abo	/e-noted.competent.ad	ult person or authorized de	clarant making his/her signature at	sove and, if applicable, the
rwo WITNESSES: (See qualificati	ions on backside.) We have witnessed the abo g an OOH-DNR by nonwritten communication	to the attending physic	ian.		oove and, if applicable, the
IWO WITNESSES: (See qualificati above-noted adult person making			ian.	darant making bis/ber signature at	oove and, if applicable, the
rwo WITNESSES: (See qualificati above noted adult person making Witness 1 signature		to the attending physic	lan. Print		sove and, if applicable, the
TWO WITNESSES: (See qualificati above-noted adult person making Witness 1 signature Vitness 2 signature	g an OOH-DNR by nonwritten communication	to the attending physic Date Date	ian. Print Prin	ed name	
TWO WITNESSES: (See qualificati bove noted adult person making Witness 1 signature Vitness 2 signature	g an OOH-DNR by nonwritten communication	to the attending physic Date Date noted person personal	ian. Print Prin	ed name	
INO WITNESSES: (See qualificati above noted adult person making Witness I signature Vitness 2 signature fotary in the State of Texas and Signature & seal:	g an OOH-DNR by nonwritten communication	to the attending physic Date Date noted person personal ed name:	ian. Print P	ed name red name signed the above noted dedaration Notary Seal	
Above-noted adult person making Witness 2 signature Witness 2 signature Kotary in the State of Texas and Signature & seal: Note: Notary cannot acknow	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mai	to the attending physic Date Date noted person personal ed name:	ian. Print Print Ily appeared before me and order in a nonwritten n	ed name ted name signed the above noted dedaration Notary Seal namer 1	n on this date:
TWO WITNESSES: (See qualificati above-noted adult person making Witness 1 signature Vitness 2 signature (otary in the State of Texas and signature & seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting.	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mai the attending physician of the above-noted p gs, including a hospital emergency departm	to the attending physic Date Date noted person personal ed name: king an OOH-DNR d erson and have noted to nent, not to initiate or	ian. Print Print Ily appeared before me and order in a nonwritten n he existence of this order in	ed name signed the above noted dedaration Notary Seal namer 1 the person's medical records. L dire	n on this date:
TWO WITNESSES: (See qualificati above-noted adult person making Witness 1 signature Vitness 2 signature (otary in the State of Texas and signature & seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting.	g an OOH-DNR by nonwritten communication I. County of The above Notary's print wiedge the witnessing of the person mail the attending physician of the above-noted p	to the attending physic Date Date noted person personal ed name: king an OOH-DNR d erson and have noted to nent, not to initiate or	ian. Print Print Ily appeared before me and order in a nonwritten n he existence of this order in	ed name signed the above noted dedaration Notary Seal namer 1 the person's medical records. L dire	n on this date:
TWO WITNESSES: (See qualificati above-noted adult person making Witness 1 signature Vitness 2 signature (otary in the State of Texas and Signature & seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital settin pacing, defibrillation, advance	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mai the attending physician of the above-noted p gs, including a hospital emergency departm	to the attending physic Date Date noted person personal ed name: king an OOH-DNR d erson and have noted to nent, not to initiate or	ian. Print Print Ily appeared before me and order in a nonwritten n he existence of this order in continue for the persons o	ed name signed the above noted dedaration Notary Seal namer 1 the person's medical records. L dire	n on this date:
WO WITNESSES: (see qualificati bove-noted adult person making Witness I signature Vitness 2 signature totary in the State of Texas and signature 8 seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting pacing, defibrillation, advance Physicaris sometime Physicaris sometime	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mai the attending physician of the above-noted p gs, including a hospital emergency depart d alrway management, artificial ventilation	to the attending physic Date Date noted person personal ed name: king an OOH-DNR erson and have noted t neent, not to initiate or te	ian. Print Print Ity appeared before me and order in a nonwrititen n bie existence of this order in continue for the persons (Date Licence #	ed name signed the above noted dedaration Notary Seal namer [the person's medical records, I dim ardiopulmonary resuscitation (C	n on this date: ect health care professiona PR), transcutaneous cardia
TWO WITNESSES: (See qualification of the second adult person making witness 1 signature Vitness 2 signature Vit	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mail the attending physician of the above-noted p gs, including a hospital emergency departs d airway management, artificial ventilation ball of the adult, who is incompetent or unable to considered inclucion or an orbita was not in the best	to the attending physic Date Date noted person personal ed name: king an OOH-DNR (reson and have noted t ment, not to initiate or b communicate and withou	ian. Print Print Ity appeared before me and order in a nonwrititen n he existence of this order in continue for the persons of Date Ucente # Tguardian, egent, proxy or re- irect health care professionals	ed name ided name isigned the above noted dedaration Notary Seal namer the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anaptal settings, incl	n on this date: ect health care professiona PR), transcutaneous cardia nonown, out (esuzeization measu ading a hospital emergency
WO WITNESSES: (See qualificati bove-noted adult person making Witness I signature Vitness 2 signature totary in the State of Texas and signature & seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting pacing, defibrillation, advance Physician's agrature Physician's agrature Physician's agrature Physician's agrature Physician's agrature Physician's agrature Physician's agrature Physician's agrature	g an OOH-DNR by nonwritten communication I County of The above Notary's print wledge the witnessing of the person mail the attending physician of the above-noted p g, including a hospital emergency depart d alrway management, artificial ventilation what of the adult, who is incompetent or unable to	to the attending physic Date Date noted person personal ed name: king an OOH-DNR (reson and have noted t ment, not to initiate or b communicate and withou	ian. Print Print Ity appeared before me and order in a nonwritten in the existence of this order in continue for the persons of Date Licema #	ed name ided name isigned the above noted dedaration Notary Seal namer the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anaptal settings, incl	n on this date: ect health care professiona PR), transcutaneous cardia nonown, out (esuzeization measu ading a hospital emergency
TWO WITNESSES: (See qualification of the second adult person making witness 1 signature Vitness 2 signature Vit	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mail the attending physician of the above-noted p gs, including a hospital emergency departs d airway management, artificial ventilation ball of the adult, who is incompetent or unable to considered inclucion or an orbita was not in the best	to the attending physic Date Date noted person personal ed name: king an OOH-DNR (reson and have noted t ment, not to initiate or b communicate and withou	ian. Print Print Ity appeared before me and order in a nonwrititen n he existence of this order in continue for the persons of Date Ucente # Tguardian, egent, proxy or re- irect health care professionals	ed name ied name isigned the above noted declaration Notary Seal namer 1 ithe person's medical records. I dire ardiopulmonary resuscitation (Cl attive: The person's specific waters are u acting in out of hospital settings, inch yanced ainway management, artificial	n on this date: ect health care professional PR), transcutaneous cardia nonown, out (esuzoitation meano ading a hospital emergency
WO WITNESSES: (See qualification of the second adult person making witness 1 signature Vitness 2 signatur	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mail the attending physician of the above-noted p gs, including a hospital emergency departs d airway management, artificial ventilation ball of the adult, who is incompetent or unable to considered inclucion or an orbita was not in the best	to the attending physic Date Date noted person personal ed name: king an OOH-DNR d erson and have noted t nent, not to initiate or be communicate and withou (niterests of the person full (CPR), transcutaneous car	ian. Print Print Print Ity appeared before me and order in a nonwritten in hie existence of this order in continue for the persons of Date License # Eguardian, agent, proxy or re- treat health care periofesionals disc paoring, proxy or re- treat health care periofesionals Printed	ed name isigned the above noted declaration isigned the above noted declaration inter Notary Seal inter 1 the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anspital settings, inclu- vanced ainway management, artificial	n on this date: ect health care professiona PR), transcutaneous cardla nonown, cut resuscitation measu uding a hospital emergency ventilation.
TWO WITNESSES: (See qualification bove-noted adult person making Witness 1 signature Witness 2 signature Kotary in the State of Texas and Signature & seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital settin pacing, defibrillation, advance Physician's agrature Witted name F. Directive by two physicians on be are, in reasonable medical judgment, department, not to initiate or contin Attending physician's agenature Signature of second physician	g an OOH-DNR by nonwritten communication I County of The above Notary's print wiedge the witnessing of the person mail the attending physician of the above-noted p gs, including a hospital emergency departs d airway management, artificial ventilation ball of the adult, who is incompetent or unable to considered inclucion or an orbita was not in the best	to the attending physic Date Date noted person personal ed name: king an OOH-DNR (erson and have noted to nent, not to initiate or b. communicate and withou interests of the person fall (CPR), transcutaneous can Date Date	Ian. Print Print Print Ity appeared before me and order in a nonwritten in he existence of this order in continue for the persons of Date License 4 Tguidlan, egent, proxy or re- treet health care professionals diac pacing, defibrillation, ad Printed Printed Printed	ed name isigned the above noted declaration isigned the above noted declaration inter Notary Seal inter 1 the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anspital settings, inclu- vanced ainway management, artificial	n on this date: ect health care professional PR), transcutaneous cardia nonown, out (esummation means ading a hospital emergency ventilation. Lic#
TWO WITNESSES: (See qualification bove-noted adult person making Witness 1 signature Witness 2 signature Vitness 3 signature Vitness 3 signature Vitness 3 signature Vitness 3 signature Vitness 4 seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting pacing, defibrillation, advance Physician's signature Vitness 1 signature Vitness 4 seal: Note: Notary cannot acknow Physician's signature Vitness 4 seal: Note: Notary cannot acknow Physician's signature Vitness 4 seal: Note: Note: Notary cannot acknow Adepartment, not to initiate or contin Attending physician's Signature of second physician Physician's decton.con digital signature	g an OOH-DNR by norwritten communication I County of The above Notary's print wledge the witnessing of the person mail the attending physician of the above-noted p g, including a hospital emergency departs d alrway management, artificial ventilation that of the solut, who is incompetent or unable to considered heriticities or are other way not in the best nue for the person: cardiopulmonary resuscitation use must meet critical illisted in Health and Salesy Code	to the attending physic Date Date noted person personal ed name: king an OOH-DNR d erson and have noted to nent, not to initiate or be communicate and withou interests of the person fail (CPR), transcutaneous car Date Date Date	ian. Print P	ed name isigned the above noted declaration isigned the above noted declaration inter Notary Seal inter 1 the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anspital settings, inclu- vanced ainway management, artificial	n on this date: ect health care professiona PR), transcutaneous cardia nonown, out (esummation means using a hospital emergency ventilation. Lic#
TWO WITNESSES: (See qualification bove-noted adult person making Witness 1 signature Witness 2 signature Vitness 3 signature Vitness 3 signature Vitness 3 signature Vitness 3 signature Vitness 4 seal: Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting pacing, defibrillation, advance Physician's signature Vitness 1 signature Vitness 4 seal: Note: Notary cannot acknow Physician's signature Vitness 4 seal: Note: Notary cannot acknow Physician's signature Vitness 4 seal: Note: Note: Notary cannot acknow Adepartment, not to initiate or contin Attending physician's Signature of second physician Physician's decton.con digital signature	g an OOH-DNR by norwritten communication County of The above The above	to the attending physic Date Date noted person personal ed name: king an OOH-DNR d erson and have noted t nent, not to initiate or her, communicate and withou interests of the person fall (CPR), transcutaneous car Date Date Date Estée (082(c) this document has been	ian. Print P	ed name isigned the above noted declaration isigned the above noted declaration inter Notary Seal inter 1 the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anspital settings, inclu- vanced ainway management, artificial	n on this date: ect health care professional PR), transcutaneous cardia nonown, out (esummation means ading a hospital emergency ventilation. Lic#
WO WITNESSES: (See qualificati bove-noted adult person making Witness 1 signature Vitness 2 signature Vitness 2 signature Vitness 2 signature Vitness 2 signature Note: Notary cannot acknow PHYSICIAN'S STATEMENT: I am acting in out-of-hospital setting pacing, defibrillation, advance Physician's signature Physician's signature Physician's signature Physician's signature Signature of second chysician on be are, in restandede medical judgments i department, not to initiate or contin Attending physician's agonature Signature of second chysician Physician's dectoration digital signature All persons who have signed at	g an OOH-DNR by nonwritten communication I County of The above Notary's print wledge the witnessing of the person mail the attending physician of the above-noted p g, including a hospital emergency departs d alrway management, artificial ventilation that of the solut, who is incompetent or unable to considered heritichie or an earth was not in the best nue for the person: cardiopulmonary resuscitation use must meet criticialisted in Health and Salesy Code	to the attending physic Date Date Date noted person personal ed name: king an OOH-DNR d erson and have noted to nent, not to initiate or herson and have noted to nent, not to initiate or be communicate and withou (CPR), transcutanous car Date Date Date Estics(082(c) this document has bee Guadian/Age	Ian. Print Print Print Print Print Print Print Print Print Date Date Printed P	ed name isigned the above noted declaration isigned the above noted declaration inter Notary Seal inter 1 the person's medical records. I dire ardiopulmonary resuscitation (Cl lative: The person's specific womes are u acting in out of Anspital settings, inclu- vanced ainway management, artificial	n on this date: ect health care professional PR), transcutaneous cardia nonown, out (esummation means ading a hospital emergency ventilation. Lic#

The Newer Version of the TDSHS Out-of-Hospital (OOH) DNR Form: Front Side

INSTRUCTIONS FOR ISSUING AN OOH-DNR ORDER

PURPOSE: The Out-of-Hospital Do-Not-Resuscitate (OOH-DNR) Order on reverse side complies with Health and Safety Code (HSC), Chapter 166 for use by qualified persons or their authorized representatives to direct health care professionals to forgo resuscitation attempts and to permit the person to have a natural death with peace and dignity. This Order does NOT affect the provision of other emergency care, including comfort care.

APPLICABILITY: This OOH-DNR Order applies to health care professionals in out-of-hospital settings, including physicians' offices, hospital clinics and emergency departments.

IMPLEMENTATION: A competent adult person, at least 18 years of age, or the person's authorized representative or qualified relative may execute or issue an OOH-DNR Order. The person's attending physician will document existence of the Order in the person's permanent medical record. The OOH-DNR Order may be executed as follows:

Section A - If an adult person is competent and at least 18 years of age, he/she will sign and date the Order in Section A.

Section B - If an adult person is incompetent or otherwise mentally or physically incapable of communication and has either a legal guardian, agent in a medical power of attorney, or proxy in a directive to physicians, the guardian, agent, or proxy may execute the OOH-DNR Order by signing and dating it in Section B. Section C - If the adult person is incompetent or otherwise mentally or physically incapable of communication and does not have a guardian, agent, or proxy, then a qualified relative may execute the OOH-DNR Order by signing and dating it in Section C.

Section D - If the person is incompetent and his/her attending physician has seen evidence of the person's previously issued proper directive to physicians or observed the person competently issue an OOH-DNR Order in a nonwritten manner, the physician may execute the Order on behalf of the person by signing and dating it in Section D.

Section E - If the person is a minor (less than 18 years of age), who has been diagnosed by a physician as suffering from a terminal or irreversible condition, then the minor's parents, legal guardian, or managing conservator may execute the OOH-DNR Order by signing and dating it in Section E.

Section F - If an adult person is incompetent or otherwise mentally or physically incapable of communication and does not have a guardian, agent, proxy, or available qualified relative to act on hisher behalf, then the attending physician may execute the OOH-DNR Order by signing and dating it in Section F with concurrence of a second physician (signing it in Section F) who is not involved in the treatment of the person or who is not a representative of the ethics or medical committee of the health care facility in which the person is a patient.

In addition, the OOH-DNR Order must be signed and dated by two competent adult witnesses, who have witnessed either the competent adult person making his/her signature in section A, or authorized declarant making his/her signature in either sections B, C, or E, and if applicable, have witnessed a competent adult person making an OOH-DNR Order by nonwritten communication to the attending physician, who must sign in Section D and also the physician's statement section. Optionally, a competent adult person or authorized declarant may sign the OOH-DNR Order in the presence of a notary public. However, a notary cannot acknowledge witnesses or notary signatures are not required when two physicians execute the OOH-DNR Order in section F. The original or a copy of a fully and properly completed OOH-DNR Order or the presence of an OOH-DNR Order and either one shall be

honored by responding health care professionals. **REVOCATION:** An OOH-DNR Order may be revoked at ANY time by the person, person's authorized representative, or physician who executed the order. Revocation can be by verbal communication to responding health care professionals, destruction of the OOH-DNR Order, or removal of all OOH-DNR identification devices from the person.

AUTOMATIC REVOCATION: An OOH-DNR Order is automatically revoked for a person known to be pregnant or in the case of unnatural or suspicious circumstances.

DEFINITIONS

Attending Physician: A physician, selected by or assigned to a person, with primary responsibility for the person's treatment and care and is licensed by the Texas Medical Board, or is properly credentialed and holds a commission in the uniformed services of the United States and is serving on active duty in this state. [HSC §166.002(12)].

Health Care Professional: Means physicians, nurses, physician assistants and emergency medical services personnel, and, unless the context requires otherwise, includes hospital emergency department personnel. [HSC §166.081(5)]

Qualified Relative: A person meeting requirements of HSC \$166.088. It states that an adult relative may execute an OOH-DNR Order on behalf of an adult person who has not executed or issued an OOH-DNR Order and is incompetent or otherwise mentally or physically incapable of communication and is without a legal guardian, agent in a medical power of attorney, or proxy in a directive to physicians, and the relative is available from one of the categories in the following priority: 1) person's spouse; 2) person's reasonably available adult children; 3) the person's parents; or, 4) the person's nearest living relative. Such qualified relative may execute an OOH-DNR Order on such described person's behalf.

Qualified Witnesses: Both witnesses must be competent adults, who have witnessed the competent adult person making his/her signature in section A, or person's authorized representatives making his/her signature in either Sections B, C, or E on the OOH-DNR Order, or if applicable, have witnessed the competent adult person making an OOH-DNR by nonwritten communication to the attending physician, who signs in Section D. Optionally, a competent adult person, guardian, agent, proxy, or qualified relative may sign the OOH-DNR Order in the presence of a notary instead of two qualified witnesses. Witnesses or notary signatures are not required when two physicians execute the order by signing Section F. One of the witnesses must meet the qualifications in HSC \$166.003(2), which requires that at least one of the witnesses not: (1) be designated by the person to make a treatment decision; (2) be related to the person by blood or marriage; (3) be entitled to any part of the person's estate after the person's death either under a will or by law; (4) have a claim at the time of the issuance of the OOH-DNR against any part of the person's estate after the person's death; or, (5) be the attending physician; (6) be an employee of the attending physician or (7) an employee of a health care facility in which the person is a patient if the employee is providing direct patient care to the patient or is an officer, director, partner, or business office employee of the health care facility.

Report problems with this form to the Texas Department of State Health Services (DSHS) or order OOH-DNR Order/forms or identification devices at (512) 834-6700.

Declarant's, Witness', Notary's, or Physician's electronic or digital signature must meet criteria outlined in HSC §166.011

Publications No. EF01-11421 - Revised July 1, 2009 by the Texas Department of State Health Services

Page 2 of 2

The Newer Version of the TDSHS Out-of-Hospital (OOH) DNR Form: Back Side

CARDIAC TRIAGE/TRANSPORT DECISION SCHEME

Patients who meet triage criteria for activation of the Regional Cardiac Plan should be transported to the <u>CLOSEST PCI CAPABLE FACILITY</u> or the <u>CLOSEST APPROPRIATE ACUTE CARE FACILITY</u> according to the plan guidelines with special consideration under the following circumstances:

- 1. If an adequate airway cannot be established and/or maintained or if the patient goes into cardiac arrest, has malignant cardiac dysrhythmias, or is acutely unstable, the patient should be transported to the <u>NEAREST</u> <u>ACUTE CARE FACILITY.</u>
- 2. Medical Control may order bypass in any situation as appropriate, such as when a facility is unable to meet hospital resource criteria, or when a patient is in need of specialty care.
- 3. If expected transport time is excessive (>25 minutes) activation of air transport should be considered.
- 4. If there is a question on whether or not to bypass a facility, on-line medical control should be contacted for the final decision.

Patients who are suspected of ACS or ECG confirmed STEMI patients should be transported to "PCI" Percutaneous Coronary Intervention Capable Facilities (Angioplasty, stent placement, or other intervention performed in the cath. lab) or the Closest Appropriate Acute Care Facility.

An Acute Care Facility may be considered appropriate if <u>all</u> of the following standards of care can be provided.

- A 12-lead ECG can be performed and interpreted within 10 minutes.
- PCI may be performed within 90 minutes or a thrombolytic can be administered within 30 minutes.

PCI is the preferred method of reperfusion if possible.

This plan is based on accepted best practice guidelines, but does allow for patient and physician preference.

A copy of the Regional Cardiac Plan as well as a current list of EMS Providers, Facility Capabilities and Designated Centers may be accessed at <u>www.spems.org</u>

STROKE TRIAGE/TRANSPORT DECISION SCHEME

Patients who meet triage criteria for activation of the SPEMS Suspected Stroke/TIA protocol should be transported to the <u>CLOSEST HIGHEST DESIGNATED STROKE FACILITY</u> or the <u>CLOSEST APPROPRIATE ACUTE CARE</u> <u>FACILITY</u> according to these protocol guidelines with special consideration under the following circumstances:

- 1. If an adequate airway cannot be established and/or maintained or if the patient goes into cardiac arrest, has malignant cardiac dysrhythmias, or is acutely unstable, the patient should be transported to the **NEAREST ACUTE CARE FACILITY.**
- 2. Medical Control may order bypass in any situation as appropriate, such as when a facility is unable to meet hospital resource criteria, or when a patient is in need of specialty care.
- 3. If expected transport time is excessive (>25 minutes) activation of air transport should be considered.
- 4. If there is a question on whether or not to bypass a facility, on-line medical control should be contacted for the final decision.

TRANSPORTATION OF SUSPECTED STROKE/TIA PATIENTS SHOULD BE BASED ON THE "TIME OF ONSET" and/or "LAST SEEN NORMAL TIME"

Historically, there has been only a single time frame or "window of opportunity" that EMS providers followed which allowed for the administration of TPA/Alteplase to those with thrombotic strokes. However, new advancements in stroke treatment have emerged that allows a surgeon to physically retrieve a thrombus from a blocked artery in the brain, also known as a Thrombectomy. The time frame or "window of opportunity" for a thrombectomy is much greater than that of TPA/Alteplase administration. Prehospital EMS providers should approach this situation differently than before by assessing the Cincinnati Stroke Scale (CSS) followed by the VAN assessment if a motor deficit is detected during the CSS. A positive VAN assessment indicates a Large Vessel Occlusion (LVO) for which a thrombectomy may be needed.

Therefore;

Patients with symptom onset times of < 4.5 hours that test **POSITIVE to CSS** and **NEGATIVE to VAN**, should be taken to the highest designated stroke facility for treatment and evaluation for interventional care.

- A non-designated facility may be appropriate if <u>all</u> the following standards of care can be provided:
 - Interpreted computed tomography (CT) imaging scan is available within 45 minutes of patient arrival
 - Thrombolytics can be administered within 60 minutes of patient arrival
 - Physician is available within 10 minutes of patient arrival

Patients with symptom onset times of < 24 hours that test **POSITIVE to CSS** and **POSITIVE to VAN** should be taken to the highest designated stroke facility capable of performing a thrombectomy.

• The EMS provider should be up to date on the current regional designated stroke facilities as well as their capabilities.

Patients with symptom onset times outside the time frames noted above should be transported to the nearest acute care facility for initial diagnosis and treatment. Similarly, a non-designated facility may be appropriate if the above standards of care can be provided.

This plan is based on accepted best practice guidelines but does allow for patient and physician preference.

PEDIATRIC TRAUMA TRIAGE/TRANSPORT DECISION SCHEME

Physiologic Criteria:

- 1. Depressed or deteriorating neurologic status (GCS ≤14) with focus on changes in the motor function
- 2. Respiratory distress or failure
- 3. Children requiring endotracheal intubation and/or ventilatory support
- 4. Shock, uncompensated or compensated
- 5. Injuries requiring any blood transfusion
- 6. Children requiring any one of the following:
 - a. Invasive monitoring (arterial and/or central venous pressure)
 - b. Intracranial pressure monitoring
 - c. Vasoactive medications

Anatomic Criteria:

- 1. Fractures and deep penetrating wounds to an extremity complicated by neurovascular or compartment injury
- 2. Fracture of two or more major long bones (such as femur, humerus)
- 3. Fracture of the axial skeleton
- 4. Spinal cord or column injuries
- 5. Traumatic amputation of an extremity with potential for replantation
- 6. Head injury when accompanied by any of the following:
 - a. Cerebrospinal fluid leaks
 - b. Open head injuries (excluding simple scalp injuries)
 - c. Depressed skull fractures
 - d. Sustained decreased level of consciousness (GCS ≤14)
 - e. Intracranial hemorrhage
- 7. Significant penetrating wounds to the head, neck, thorax, abdomen or pelvis including the groin
- 8. Pelvic fracture
- 9. Significant blunt injury to the chest, abdomen or neck (e.g. hanging or clothesline MOI's)

Other Criteria:

- 1. Suspicion for Child Maltreatment as evidenced by:
 - a. Injuries sustained with no reported explanation
 - b. Injuries sustained that do not match the developmental capability of the patient
 - c. History of apparent life threatening event
 - d. Upper extremity fractures in a non-ambulatory child

TRIAGE/TRANSPORT GUIDE:

• Critically injured pediatric patients meeting the above criteria should be taken to the closest verified Pediatric

Trauma Center and if unavailable the closest appropriate Adult Trauma Center with Pediatric capabilities

• If immediate interventions are required, transport to the closest appropriate acute care facility

The full BRAC Regional Pediatric Plan can be viewed at www.b-rac.org

PEDIATRIC BURN TRIAGE/TRANSPORT DECISION SCHEME

American Burn Association Triage/Transfer Criteria:

A burn center may treat adults, children, or both. Burn injuries that should be referred to a burn center include the following:

- 1. Partial-thickness burns of greater than 10 percent of the total body surface area.
- 2. Burns that involve the face, hands, feet, genitalia, perineum, or major joints.
- 3. Third-degree burns in any age group.
- 4. Electrical burns, including lightning injury.
- 5. Chemical burns.
- 6. Inhalation injury.
- 7. Burn injury in patients with preexisting medical disorders that could complicate management, prolong recovery, or affect mortality.
- 8. Physician judgment will be necessary in such situations and should be in concert with the regional medical control plan and triage protocols.
- 9. Burn injury in patients who will require special social, emotional, or rehabilitative intervention.

TRIAGE/TRANSPORT GUIDE:

Burns in Children:

<u>Children with burns should be transferred to a burn center verified to treat children. In the absence</u> of a regional pediatric burn center, an adult burn center may serve as a second option for the management of pediatric burns.

Burns and Trauma in Children:

Burns and concomitant trauma (such as fractures) when the burn injury poses the greatest risk of morbidity or mortality.

If the trauma poses the greater immediate risk, the patient's condition may be stabilized initially in a Pediatric Trauma Center before transfer to a burn center.

Other Criteria for Transport:

- 1. Children requiring pediatric intensive care other than for close observation.
- 2. Any child who may benefit from consultation with, or transport to, a Pediatric Trauma Center or a Pediatric Intensive Care Unit.
- 3. Children with injuries suspicious of child maltreatment e.g. inflicted burn injury

Reference: Resources for the Optimal Care of the Injured Patient: 2014

The full BRAC Regional Pediatric Plan can be viewed at www.b-rac.org

PEDIATRIC NON-TRAUMA TRIAGE/TRANSPORT DECISION SCHEME

Physiologic Criteria

- 1. Depressed or deteriorating neurologic status (GCS≤14).
- 2. Severe respiratory distress and/or respiratory failure
- 3. Children requiring endotracheal intubation and/or ventilatory support.
- 4. Serious cardiac rhythm disturbances,
- 5. Status post cardiopulmonary arrest.
- 6. Heart failure.
- 7. Shock responding inadequately to fluid resuscitation.
- 8. Children requiring any one of the following:
 - a. Arterial pressure monitoring.
 - b. Central venous pressure or pulmonary artery monitoring.
 - c. Intracranial pressure monitoring.
 - d. Vasoactive medications.
 - e. Treatment for severe hypothermia or hyperthermia
 - f. Treatment for hepatic failure.
 - g. Treatment for renal failure, acute or chronic requiring immediate dialysis.

Other Criteria

- 1. Near drowning with any history of loss of consciousness, unstable vital signs or respiratory problems.
- 2. Status epilepticus.
- 3. Potentially dangerous envenomation. Use of a snakebite protocol is encouraged
- 4. Potentially life threatening ingestion of, or exposure to, a toxic substance.
- 5. Severe electrolyte imbalances.
- 6. Severe metabolic disturbances.
- 7. Severe dehydration.
- 8. Potentially life-threatening infections, including sepsis.
- 9. Children requiring intensive care other than for close observation.
- 10. Any child who may benefit from consultation with, or transfer to, a Pediatric Intensive Care Unit
- 11. Suspicion for child maltreatment. e.g. found "down" for no apparent reason
- 12. Any condition that exceeds the capability of the facility

TRIAGE/TRANSPORT GUIDE:

• Critically ill pediatric patients meeting the above criteria should be taken to the closest Children's Hospital or

hospital with Pediatric Critical Care capabilities

• If immediate interventions are required, transport to the closest appropriate acute care facility

The full BRAC Regional Pediatric Plan can be viewed at www.b-rac.org

TRAUMA TRIAGE CRITERIA FOR EMS FACILITY BYPASS AND TRANSFER

Patients who meet triage criteria for activation of the regional trauma system plan will be transported <u>DIRECTLY TO</u> <u>AN APPROPRIATE TRAUMA FACILITY</u>, rather than to the nearest hospital, <u>EXCEPT</u> under the following circumstances:

- 1. If an adequate airway cannot be established and/or maintained, or in cases of traumatic cardiac arrest, the patient should be taken to the **NEAREST ACUTE CARE FACILITY** for stabilization.
- 2.A Level III (Basic) facility may be appropriate for immediate evaluation and stabilization if the expected transport time to a Level I or Level II facility is excessive (>25 minutes).
- 3.Medical Control may order a bypass in any of the above situations as appropriate, such as when a facility is unable to meet hospital resource criteria, or when the patient is in need of specialty care.
- 4.If expected transport time is excessive (>25 minutes), or if expected extrication time is lengthy (>15 minutes), activation of air transport resources should be considered.
- 5. If there is a question on whether or not to bypass a facility, on-line medical control should be consulted for the final decision.

CRITICAL PATIENTS

Should be evaluated at a Level I or Level II Trauma Center. Patients who are physiologically and hemodynamically unstable whose injuries may include:

- Chest:
 - Major chest wall injury
 - Penetrating thoracic wound
- Pelvis:
 - Pelvic ring disruption with shock requiring >5 units transfusion
- Abdomen:
 - Blunt trauma with hypotension
 - Penetrating abdominal wound
- Multiple system trauma:
 - Chest injury with head injury
 - Abdominal or pelvic injury with head injury
- Specialized problems:
 - 2° or 3° burns >10% TBSA or involving airway
 - Barotrauma
 - Uncontrolled hemorrhage
 - 2nd/3rd trimester pregnancy

URGENT PATIENTS

Should be evaluated at a Level I or Level II Trauma Center. Patients who are physiologically and hemodynamically unstable whose injuries may include:

- Central nervous system:
 - Prolonged loss of consciousness, posturing,

paralysis or lateralizing sign

- Spinal injuries with or without deficit
- Glasgow Coma Score (GCS) <10
- Open, penetrating or depressed skull fracture
- CSF leak
- GCS deterioration ≥2
- Thoracic:
 - Suspected cardiac/great vessel injury
 - Possible requirement for prolonged mechanical ventilation
 - Respiratory distress with rate >35 or <10
- Abdomen:
 - Blunt trauma without hypotension
- Multiple system trauma:
 - Severe facial injury with head injury
 - Specialized problems:
 - Carbon monoxide poisoning
 - Severe maxillofacial or neck injuries
 - Revised Trauma Score (RTS) ≤11
 - Open fractures
- Secondary deterioration (Late Sequelae):
 - Patients requiring mechanical ventilation
 - Sepsis
 - Organ system(s) failure
 - Osteomyelitis

Continued on Next Page

TRAUMA TRIAGE CRITERIA FOR EMS FACILITY BYPASS AND TRANSFER

(Continued)

CATEGORY III PATIENTS

May be evaluated at a Level III Trauma Center. Patients who are physiologically and hemodynamically stable whose injuries may include:

- Central nervous system:
 - Transient loss of consciousness
- Chest:
 - Injuries not producing respiratory distress
 - Rib fractures without flail segments
- Abdomen:
 - Blunt trauma without hypotension
- Specialized problems:
 - Closed fractures
 - · Soft tissue injuries with controlled hemorrhage
 - 2nd/3rd trimester pregnancy

CATEGORY IV PATIENTS

May be evaluated at an appropriate trauma facility. Patients who are continually normotensive and/or hemodynamically stable, but whose injuries may include:

- Specialized problems:
 - · Closed fractures without neurological deficit
 - Moderate soft tissue injuries

GLASGOW COMA SCORE & REVISED TRAUMA SCORE

To calculate the Revised Trauma Score:

- 1. Calculate the Glasgow Coma Score
- 2. Determine score components based on Glasgow Coma Score, Respiratory Rate, and Systolic Blood Pressure.
- 3. Add score components to determine REVISED TRAUMA SCORE.

GLASGOW COMA SCORE - Ad	ult & Child	
MOTOR RESPONSE	VERBAL RESPONSE	EYE RESPONSE
1-No Response	1-No Response	1-No Response
2-Abnormal Extension	2-Incomprehensible Sounds	2-To Pain
3-Abnormal Flexion	3-Inappropraite Words	3-To Verbal Command
4-Withdrawal	4-Confused/Disoriented	4-Spontaneous
5-Localizes Pain	5-Oriented	
6-Obeys Command		
-		
GLASGOW COMA SCORE – Ch	ild & Infant	

ASGOW COMA SCORE – Child & Infant

MOTOR RESPONSE	VERBAL RESPONSE	EYE RESPONSE
1-No Response	1-No Response	1-No Response
2-Abnormal Extension	2-Moans, Grunts	2-To Pain
3-Abnormal Flexion	3-Cries to Pain	3-To Speech
4-Withdraws to Pain	4-Irritable Cries	4-Spontaneous
5-Localizes Pain	5-Coos, Babbles	
6-Spontaneous		

REVISED TRAUMA SCORE – Adult & Child

GLASGOW COMA SCORE	RESPIRATORY RATE	SYSTOLIC BLOOD PRESSURE
0=3	0=0	0=0
1=4-5	1=1-5	1=1-49
2=6-8	2=6-9	2=50-75
3=9-12	3=>29	3=76-89
4=13-15	4=10-29	4=>89

Score: 0-12 (decreasing with increasing injury severity) Patients with Revised Trauma Score of 11 or less require care at a Level I or Level II Trauma Center.

REVISED TRAUMA SCORE – Child & Infant

Score	Weight	Airway	BP*	Level of Consciousness	Open Wound	Fractures
+2	>20kg (44lbs)	Normal	>90mmHg	Awake	None	None
+1	10-20kg (22-44lbs)	Maintainable with O ₂	50-90mmHg	Obtunded or any LOC	Minor	Closed Fracture
-1	<10kg (22lbs)	Intubated	<50mmHg	Comatose	Major or Penetrating	Open or Multiple

*In the absence of a Blood Pressure reading, the BP may be estimated by the point at which a pulse is palpable as follows: +2 - Brachial, +1 - Groin, -1 - No Pulse Palpable

Score: 0-12 (decreasing with increasing injury severity) Patients with Revised Trauma Score of 11 or less require care at a Level I or Level II Trauma Center.

GUIDELINES FOR TRAUMA TEAM ACTIVATION

Adult Patients (≥16 years of age)

Level I

- 1. Unstable vital signs:
 - Confirmed Systolic BP <90mmHg at any time
 - Sustained Pulse <50 or >120
- 2. Respiratory compromise/obstruction and or intubation (pre-hospital)
 - Respiratory rate <10 or >35 per minute
 - Unsecured airway
 - •Clinical symptoms of hypoxia
- 3. Glasgow Coma Scale < 8, with mechanism related to trauma
- 4. Gun shot wound to abdomen, neck, or chest
- 5. RTS ≤10 on arrival
- 6. Severe multi-system trauma
- 7. Traumatic amputation of limb (with clinical instability or associated injuries)
- 8. Transfer patients requiring fluid, pressors, or blood to maintain vital signs
- 9. Burn injuries
 - 50% TBSA 2° & 3° burns (all ages)
 - High voltage electrical burns with cardiac arrhythmias or significant tissue damage,
 - Inhalation injuries with respiratory distress
- 10. EC physician discretion

Level II

- 1. Intubated patients transferred from another facility
- 2. Flail Chest
- 3. Trauma resulting in an open long bone fracture
- 4. Pelvic fracture
- 5. Penetrating injury to extremities and stab wounds to the trunk
- 6. MVC:
 - Un-restrained rollover
 - Ejection from the vehicle
- 7. MCC:
 - No Helmet
 - MCC traveling > 20 mph
- 8. Pedestrians struck by a vehicle moving >20mph
- 9. Glasgow Coma Scale >8 but <13, with mechanism related to trauma
- 10. Falls >20 feet
- 11. Burns:
 - > 10% TBSA 2° or 3° burn <10 or > 50 years of age
 - > 20% but < 50% TBSA 2° (all ages)
 - > 20% TBSA 3° (all ages)
 - All other electrical burns
 - All chemical burns
- 12. Symptomatic Carbon monoxide (CO) poisoning
- 13. EC physician discretion

Level III

- 1. Traumatically injured patients not otherwise defined
- 2. Clinically stable patients with injuries identified after EC work-up
- 3. Injured patients requiring subspecialty consult
- 4. Burns not otherwise defined

GUIDELINES FOR TRAUMA TEAM ACTIVATION (continued)

Pediatric Patients (< 16 years of age)

Level I

1. Unstable vital signs:

Age	Systolic BP	Pulse Rate	Respiratory Rate
Birth – 1 year	Capillary Refill >4 seconds	<80 or >180	>60
1 – 5 years	<70mmHg	<60 or >160	>50
6 – 14 years	<80mmHg	<50 or >140	>50

- 2. Respiratory compromise/obstruction and or intubation (pre-hospital)
 - Unsecured airway
 - Clinical symptoms of hypoxia
- 3. Glasgow Coma Scale < 8, with mechanism related to trauma
- 4. Any Gun shot wound
- 5. Any penetrating trauma to the torso (chest or abdomen)
- 6. RTS ≤ 10 on arrival
- 7. Severe multi-system trauma
- 8. Traumatic amputation of limb (with clinical instability or associated injuries)
- 9. Transfer patients requiring fluid, pressors, or blood to maintain vital signs
- 10. Burn injuries
 - 50% TBSA 2° & 3° burns (all ages)
 - High voltage electrical burns with cardiac arrhythmias or significant tissue damage,
 - Inhalation injuries with respiratory distress
- 11. EC physician discretion

Level II

- 1. Intubated patients transferred from another facility
- 2. Flail Chest
- 3. Trauma resulting in an open long bone fracture
- 4. Pelvic fracture
- 5. Penetrating injury to extremities
- 6. MVC:
 - Un-restrained rollover
 - Ejection from the vehicle
- 7. MCC:
 - No Helmet
 - MCC traveling > 20 mph
- 8. Child struck or run over by a motor vehicle or trailer
- 9. Near drowning
- 10. Glasgow Coma Scale >8 but <13, with mechanism related to trauma
- 11. Falls >20 feet
- 12. Burns:
 - > 10% TBSA 2° or 3° burn <10 or > 50 years of age
 - > 20% but < 50% TBSA 2° (all ages)
 - >20% TBSA 3° (all ages)
 - All other electrical burns
 - All chemical burns
- 13. Symptomatic Carbon monoxide (CO) poisoning
- 14. EC physician discretion

Level III

- 1. Traumatically injured patients not otherwise defined
- 2. Clinically stable patients with injuries identified after EC work-up
- 3. Injured patients requiring subspecialty consult
- 4. Burns not otherwise defined





CRITERIA FOR THE CONSIDERATION OF AIR MEDICAL TRANSPORT FOR TRAUMA PATIENTS

- Lengthy extrication of the patient at the scene and the severity of the patient's injuries require delivery of a critical care team to the scene.
- One or more of the following mechanisms of injury with a motor vehicle collision present:
 - There had been structural intrusion into the patient's space in the vehicle;
 - The patient was ejected from the vehicle;
 - Another person in the same vehicle died;
 - The patient was a pedestrian struck by a vehicle traveling more than 20mph;
 - The patient was not wearing a safety belt in a car which was overturned;
 - The patient was thrown from a motorcycle traveling more than 20mph.
- The front bumper of the vehicle was displaced to the rear by more than 30 inches, or the front axle was displaced to the rear.
- The patient fell from a height of greater than 20 feet.
- The patient experienced a penetrating injury between the mid-thigh and the head.
- The patient experienced an amputation, or near amputation, and required timely evaluation for possible reimplantation.
- The patient experienced a scalping or degloving injury.
- The patient experienced a severe hemorrhage. Included are those patients with a systolic blood pressure of less than 90mmHg after initial volume resuscitation and those requiring ongoing blood transfusions to maintain a stable blood pressure.
- The patient experienced 2°/3° degree burns of the skin greater than 15 percent of the body surface, or major burns of the face, hands, feet, or perineum, or associated with an airway or inhalation injury.
- The patient experienced, or had great potential to experience, injury to the spinal cord, spinal column, or neurologic deficit.
- The patient suffered injuries to the face or neck which might result in an unstable or potentially unstable airway and might require invasive procedures (such as endotracheal or nasotracheal intubation, tracheostomy cricothyrotomy) to stabilize the airway.
- The patient had a score from an objective ranking system for trauma (such as the Trauma Score, Revised Trauma Score, CRAMS, Glasgow Coma Scale, etc.) at the scene or at the referring hospital's emergency department which indicated a severe injury.
- The patient is a child less than five years of age with multiple traumatic injuries.
- The patient is greater than 55 years of age and has multiple traumatic injuries, whether with or without preexisting illness, such as diabetes mellitus, coronary artery disease, chronic obstructive lung disease, or chronic renal failure.
- The patient is an adult with respiratory rate of less than 10 or greater than 35 breaths per minute, or a heart rate of less than 60 or greater than 120 beats per minute.

Source: FLIGHT NURSING: PRACTICE AND PRINCIPLES, 1991

SOUTH PLAINS EMS EQUIPMENT LIST

BLS UNITS

- 1- SAED with 2 sets of defibrillator pads/paddles to accommodate adult patients and 2 sets of defibrillator pads/paddles to accommodate pediatric patients. However, if the BLS unit already stocks a monitor/defibrillator/SAED another SAED is not required. (If the SAED stocked does not support pediatric defibrillation a variance must be filled out through the SPEMS office and then pediatric defibrillation pads are not mandatory. The Variance must be signed by the Medical Director and a copy placed in each set of protocols). A charged spare battery must accompany the unit as well as the one powering the unit. However, an alternative power source may take the place of the spare battery. (SAED with sealed 5 year batteries need not to have a spare)
- 1- Portable suction (no foot pump or bulb type) with charged spare battery if the unit is battery powered. However, an alternative power source may take the place of the spare battery.
- 1- Vehicle mounted suction
- 3ea-suction tubing, rigid suction tips, and suction canisters
- 1ea- Bag valve mask (adult and child sizes)
- 1- ResQPOD (Optional)
- 1ea- Oral airway (#2 through #6)
- lea Nasal airway (20fr through 36fr)
- 3ea- Adult nonrebreather masks, pediatric nonrebreather masks, adult handheld nebulizers or nebulizer masks, adult nasal cannulas, and oxygen tubings. **Note**: if nonrebreather masks or nebulizers contain removable oxygen tubing, then the requirement for oxygen tubing is met
- 2- Portable oxygen cylinders
- 1- Portable oxygen regulator
- 1- Piped-in oxygen with regulator (M or H cylinders)
- 1- Pulse Oximeter device with charged spare batteries
- 2ea- C-collars (to accommodate adult, child, and infant)
- 1- Extremity splint for all extremities
- 1ea- Traction splint (adult, child sizes)
- 2- Full immobilization devices with straps
- 1- Short board or KED-type extrication device (does not have to be a KED brand)
- 1ea- Blood pressure cuffs (adult, child sizes)
- 1- Stethoscope
- 1- Glucometer
- 2- Appropriate glucometer test strips
- 2- Lancet/needle
- 3ea-1cc syringes, 3cc syringes, and 10cc syringes
- 1- Nasal-Mucosal Atomization Device (MAD)
- 3- Hypodermic needles (if IM Epinephrine is stocked) (sizes to be determined by EMS provider's needs)
- 2- Multi trauma dressings
- 25- START Triage Tags (START = Simple Triage and Rapid Transport)
- 1- QuickClot (must be heat free formula)
- 1- Commercial Tourniquet Device
- 24- Sterile gauze pads
- 5- Soft roller adhering bandages

03/01/2022

- 2- Rolls of adhesive tape
- 2- Triangular bandages
- 2- Sterile burn sheets
- 3- Vaseline gauze
- 10- Alcohol preps
- 1- Bandage shears
- 1- Pen light
- 1- Multilevel stretcher with all patient securing straps as recommended by the stretcher's manufacturer and at least 2 sets of clean sheets and blankets
- 1- Mast pants (Optional)
- 1ea- King LT-D or LTS-D airway sizes 0, 1, 2, & 2.5
- 1ea- King LTS-D airway sizes 3, 4, & 5
- 1- Puncture resistant sharps container
- 1- Emergency Response Guidebook, most current edition
- 1set- Emergency warning devices (at least 3 emergency triangles)
- 10- Protective, non-porous gloves
- 2- Medical eye protection
- 5- N-95 or greater protective Masks
- 2- Medical protective gowns or equivalent
- 2- Personal cleansing supplies
- 5- Biohazard bags
- 1- Working flashlight
- 1- Fire extinguisher
- 1- No smoking signs in the cab and patient compartment
- 1- Thermometer (may be oral, tympanic, or skin monitoring)
- 1- Mechanical CPR Compression Device (Optional). If carried, services are to follow manufacturer's recommendation assuring compliance with current AHA CPR guidelines
- 4- Cold packs or equivalent

BLS NEONATAL EQUIPMENT

- 1- Sealed OB kit with non-porous infant insulator, umbilical cord clamps and bulb aspirator
- 1-Broselow Pediatric Emergency Tape or equivalent
- 1-Infant Bag valve mask (BVM)
- 1-#1 Oral airway

Note: much of the other equipment, supplies, and drugs listed in other sections of the Equipment List are also designed to be utilized for neonatal care and this section does not exclude the use of any equipment, supplies, or drugs for neonatal patients as allowed by these treatment Protocols.

BLS MEDICATIONS

- 1- Activated Charcoal, 50g
- 20- Acetaminophen, 500mg tablets
- 10- Aspirin, 325mg tablets
- 5- Duo-Neb: 0.5mg Ipratropium Bromide and 3mg of Albuterol Sulfate in 3ml
- 1- Epinephrine Auto-Injector (Adult) 0.3mg/unit
- 1- Epinephrine Auto-Injector (Pediatric) 0.15mg/unit
- 2- Epinephrine (1:1,000) 1mg/1cc (if stocked at the BLS level, appropriate training required)
- 20- Ibuprofen, 200mg tablets

03/01/2022

- 1- Glucagon 1mg/unit (Optional)
- 2- Levalbuterol (Xopenex) 1.25mg/3ml (Optional)
- 1- Liquid Children's Motrin 100mg/5ml
- 2- Naloxone (Narcan) 2mg/2cc
- 1- bottle- Nitroglycerin 0.4mg tablet or spray
- 1- Oral Glucose 15g Tube
- Oxygen

Services under SPEMS medical direction may carry Epinephrine Auto-Injectors to accommodate both adult and pediatric patients **AND/OR** Epinephrine (1:1,000) 1mg/1cc. However, Epinephrine (1:1,000) can only be carried if all active ECA's, EMT's and Advanced EMTs are appropriately trained on IM injections (and the standing Allergic Reaction Protocol). This training must be documented including location, date, and time. Documentation must be readily accessible upon inspection.

ALS & ALS-CAPABLE UNITS

ALL BLS EQUIPMENT AND:

- Advanced airway equipment including equipment required to perform endotracheal intubation including but not limited to:
 - 1-Laryngoscope handle with appropriate spare batteries
 - 1ea- Laryngoscope blades (Miller 1,2,3,4 and Mac 1,2,3,4)
 - 1ea- Extra laryngoscope bulbs (small and large)(extra bulbs not required for fiber optic laryngoscope sets)
 - 2ea- Endotracheal tubes (adult, pediatric sizes, 4 thru 9.0)
 - ♦ 1- ET tube holder
 - 1ea- Stylet (adult, pediatric sizes)
- 1ea- ET tube introducer (i.e. Bougie) (adult and pediatric)
- 2- Soft suction catheters for suctioning of ET tube
- 1- Carbon Dioxide monitor or detector (if a Carbon Dioxide detector (i.e. Easy Cap) is solely utilized an adult and pediatric must be stocked)
- 1ea- Magill forceps (adult and pediatric)
- Equipment & supplies to establish intravenous infusions for adult & pediatric patients to include but not limited to:
 - ◆ 5ea- IV catheters (16ga, 18ga, 20ga, 22ga)
 - ♦ 4-1000cc Lactated Ringers or equivalent volume
 - ♦ 4-1000cc Normal Saline or equivalent volume
 - ♦ 4-Macro IV tubing
 - 2-Micro IV tubing (60 drop set)
 (If dial-a-flow tubing is used then a total of 6 IV tubes are required)
 - 4-Securing devices (nonporous tape or Veni-guard devices are acceptable)
- 1- EZ IO Driver with spare batteries (EZ IO Drivers with the non-replaceable battery need not have a spare)
- 1- EZ IO PD needle
- 1- Adult EZ-IO needle
- 1- EZ-IO LD needle (optional)
- 1- EZ-STABILIZER (optional)
- 2ea- Blue, Red, Purple, and Green blood specimen tubes or their equivalent
- 2- Gastric tubes
- 2-14ga X 2" or longer IV catheters for pleural decompression

- 1- Morgan Lenses (Optional)
- 1- Copy of current, signed SPEMS treatment protocols for Advanced EMT

ALS NEONATAL EQUIPMENT

In addition to the BLS Neonatal Equipment list, all ALS or ALS Capable units must be equipped with:

- 1- Miller size 0 laryngoscope blade with extra bulb (extra bulb not required for fiber optic laryngoscope blade)
- 2ea- Endotracheal tubes (sizes, 2.5, 3.0, and 3.5)
- 5-24ga IV catheters

Note: much of the other equipment, supplies, and drugs listed in other sections of the Equipment List are also designed to be utilized for neonatal care and this section does not exclude the use of any equipment, supplies, or drugs for neonatal patients as allowed by these treatment Protocols.

ALS MEDICATIONS

- 2- Dextrose 10% (D₁₀W), 250cc prefilled IV solution
 - **NOTE**: Until current supplies of Dextrose 50% (D₅₀W), 25g/50cc are exhausted or expires, EMS services can meet the requirement for Dextrose 10% by carrying 2-Dextrose 50% (D₅₀W), 25g/50cc AND 2- 250cc bags of NS. D₁₀W can then be achieved by removal of 50cc of the NS and injecting the 50cc of D₅₀W into the IV bag. Once current supplies of D₅₀W are exhausted, EMS services are required to stock the D₁₀W premixed solution in a 250cc bag.
- 1- Diphenhydramine (Benadryl), 50mg/cc
- 4- Epinephrine (1:10,000), 1mg/10cc
- 1- Lidocaine (Xylocaine), 100mg/5cc
- 2- Ondansetron (Zofran) 4mg/2cc
- 1- Racemic Epinephrine 2.25% 11.25mg/0.5ml (optional)
- 2- IV Acetaminophen (1,000mg/100cc) Pre-mixed IV solution. If premixed is not available, services may carry 2 vials of IV Acetaminophen of 1,000mg and 2 100cc bags of 0.9% normal saline (NS) or D5W
- If you have medical direction for any medications or invasive equipment not listed here, you must attach written authorization for the use of such. This document must be signed by the SPEMS Medical Director. However, non-invasive equipment (example: Vacu-Mattress, vein finder, thermometer, etc.) does not require written authorization by medical direction or additions to the equipment list.
- All of the services under my medical direction must carry the minimum amount of equipment and medications listed above, and may carry more according to their run demand and patient care needs.

Ca

SPEMS Medical Director

Date 03/01/2022

Service Director

Date 03/01/2022



ADVANCED EMT

TRAUMATIC EMERGENCIES



03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

Page 1







ADVANCED EMT

RESPIRATORY EMERGENCIES



NEAR DROWNING





• Consider spinal cord trauma, air embolism, hypothermia, alcohol or drug ingestion, hypoglycemia, seizures and myocardial infarction as accompanying problems or underlying causes.

All near drowning patients should be transported for observation & evaluation, no matter how mild the episode appears to be.

• Air transport should be considered to expedite the patient's arrival at the hospital.

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

RESPIRATORY DISTRESS







ADVANCED EMT

CARDIOVASCULAR EMERGENCIES

CARDIAC ARREST/SEMIAUTOMATIC EXTERNAL DEFIBRILLATION*





** Ideally chest compressions should be interrupted only for rhythm checks and actual defibrillations. The Current AHA guidelines state that when CPR is indicated the provider should perform 5 cycles (2 Minutes) of chest compressions. Continue CPR while the defibrillator is charging. Providers must organize care to ensure that chest compressions, initial and subsequent defibrillations are not delayed due to placement of advanced airways or to obtain vascular access.

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.


CARDIOGENIC SHOCK





PEDIATRIC

Fluid challenge 20cc/kg over 10 minutes. Repeat until clinical signs of adequate perfusion are present. Monitor patient for pulmonary edema.

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.







See Additional Information on Next Page

LVAD (Continued)



LVAD stands for Left Ventricular Assist Device. It is an electromechanical device for assisting cardiac circulation that is designed to assist a failing heart. Patients may or may not be on a heart transplant list. These patients will most likely have an internal pacemaker and an internal defibrillation device.

The LVAD is implanted into the left upper abdominal quadrant and produces a humming sound when functioning. It is critical that the pump be auscultated to assure proper functioning. In the event of arrest, the LVAD device should produce sufficient blood flow without the performance of chest compressions. Chest compressions can cause dislodgement of the device and could result in death from blood loss.

Key Notes for caring for LVAD Patients:

- Always talk to family/caregivers since they will have specialized training and knowledge about the device. An appropriate knowledgeable/trained family member should accompany the patient to the hospital.
- CALL THE VAD COORDINATOR EARLY per patient/family instructions or as listed on the device. The VAD coordinators are available 24/7 and should be an integral part of the treatment plan.
- Questions to ask: Does the patient have a DNR? Can the patient be cardioverted or defibrillated if needed?
- Chest compressions should NOT be given unless the pump has stopped working (not humming).
- Common complication in LVAD patients include stroke and TIA (incidence up to 25%), bleeding, dysrhythmias, and infection.
- The cardiac monitor and 12-Lead ECG are not affected by the LVAD and will provide important information.
- LVAD patients are pre-load dependent. Consider that a FLUID BOLUS can often reverse hypoperfusion.
- Transport patients with ALL device equipment including any instructions, hand pumps, backup batteries, primary and secondary controllers, as well as knowledgeable family members or caregivers. Most patients will have a "Go Bag" available. A charged spare battery should be part of the "Go Bag".
- Remember that it is unlikely an auscultated or palpated blood pressure can be measured. Pulse oximetry may be inaccurate.
- If patient is unconscious with no respiratory effort and the ECG shows V-Fib or V-Tach, defibrillation is necessary
- CPAP CANNOT be used on patients with LVAD

Prehospital Medications to Avoid with LVAD Patients:

Anti-arrhythmics May be used only if patient is in cardiac arrest	
Amiodarone	Amiodarone and Lidocaine may be used in cardiac
Diltiazem (Cardizem) and other calcium channel	arrest. Calcium channel blockers should not be
blockers	administered due to the negative inotropic affect and
Lidocaine	potential to develop hypotension.
Anti-Coagulants/ Anti-platelet properties	
Aspirin	Patient is already on anti-coagulants and adding aspirin
	may increase bleeding risks or delay healing
Anti-hypertensives	
Labetalol (Normodyne)	Peripheral vasodilation or significant reduction in
	blood pressure may decrease preload. The LVAD
	patient is dependent on preload for optimal pump
	function.
Benzodiazepines	
Diazepam (Valium)	May result in peripheral vasodilation that can reduce
Lorazepam (Ativan)	preload to the heart. For continuous seizures, use
Midazolam (Versed)	lowest effective dose.
Elementals	
Magnesium Sulfate	Risk for developing hypotension and dysrhythmias
Sympathomimetic Medications May be used only if patient is in cardiac arrest	
Epinephrine 1:10,000	Drugs that stimulate the sympathetic response may
Dopamine	increase myocardial oxygen demand
Repeated doses of nebulized beta-agonists	Increases myocardial oxygen demands
Vasodilatory Medications	
Nitroglycerin	Peripheral vasodilation will decrease preload. The
Morphine	LVAD patient is dependent on preload



Fluid challenge 20cc/kg over 10 minutes. Repeat until clinical signs of adequate perfusion are present. Monitor patient for pulmonary edema.

If patient's condition changes, refer to appropriate algorithm.

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.



ADVANCED EMT

OTHER MEDICAL EMERGENCIES

ALLERGIC REACTION*





PEDIATRIC DOSE Epinephrine, 0.15mg, IM, Using Auto-Injector or Epinephrine, (1:1,000), 0.01mg/kg to a max of 0.15mg (0.15cc), IM, if properly trained Duo-Neb (0.5mg Ipratropium Bromide with 3mg Albuterol), by nebulizer, dose and frequency same as adult Benadryl 1mg/kg, IV to max of 50mg Xopenex (optional), 1.25mg via nebulizer, dose and frequency same as adult

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.



03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.





03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

HAZARDOUS/TOXIC MATERIAL EXPOSURE



- 1. Observe Hazmat Precautions*
- 2. Do Not Enter Incident Area Without Appropriate Protective Clothing/Respiratory Equipment
- 3. Evacuate Patients From Exposure Without Risking EMS Personnel Safety
- 4. In Cooperation With Police/Fire Authorities, Evacuate/Isolate Scene
- 5. Attempt to Identify Nature of Hazardous Material as Soon as Possible
- 6. Request Paramedic Backup



03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.







Suspect Heat Stroke in any Patient with an Altered Level of Consciousness in a Hot Environment

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

HYPOVOLEMIA* (NON-TRAUMATIC)





PEDIATRIC

Fluid challenge 20cc/kg over 10 minutes. Repeat until clinical signs of adequate perfusion are present. Monitor patient for pulmonary edema.

****POSITIVE TILT TEST**

Pulse Rate Increases by 20, Systolic BP Decreases by 20mmHg, or Diastolic BP Decreases by 10mmHg when Patient is Raised from Supine to Sitting position **OR** Patient will Not Tolerate Being Raised From Supine to Sitting Position Because of Weakness, Dizziness, Presyncope, or Syncope.

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

NEONATAL RESUSCITATION





* Ideally chest compressions should be interrupted only for rhythm checks and actual defibrillations. The Current AHA guidelines state that when CPR is indicated the provider should perform 5 cycles (2 Minutes) of chest compressions. Continue CPR while the defibrillator is charging. Providers must organize care to ensure that chest compressions, initial and subsequent defibrillations are not delayed due to placement of advanced airways or to obtain vascular access.

03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

OBSTETRIC EMERGENCY





03/01/2022

IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

POISONING/OVERDOSE





03/01/2022





- 1. Remove Patient From Potentially Harmful Environment; Do NOT Forcefully Restrain Patient
- 2. Secure Airway
- 3. Oxygen
- 4. Assist Ventilations if Respirations Inadequate (Insert OPA/NPA)
- 5. IV, **NS**, TKO
- 6. Determine Blood Glucose





**

PEDIATRIC DOSE

- •Dextrose 10% (25g/250cc), Wide-open with a 60gtts IV set until pateint becomes responsive. Once responsive obtain a BGL. If BGL is ≥ 90mg/dL slow D10 infusion to a TKO rate and monitor to maintain desired effect. Monitor Lung sounds closely
- ***•Liquid Motrin (Ibuprofen), 10mg/kg, PO, to a max of 800mg per single dose Post seizures with temperature > 100.4 and adequate LOC.

Glucagon, 0.5mg, IM (Optional)

- Dextrose 10%, 25g/250cc, should be infused at a wide-open rate until the patient becomes responsive. Once responsive obtain BGL. If BGL <u>></u> 90mg/dL slow D10 infusion to a TKO rate and monitor to maintain desired effect.
- If venous access is unobtainable administer Glucagon, 1mg/unit, IM (Optional). Appropriate training and testing must be documented prior to administration of Glucagon via IM injection. Do Not establish EZ-IO unless BGL is < 50 mg/dL and/or pt condition warrants



IF PATIENT'S CONDITION IS UNSTABLE OR DETERIORATES DURING TRANSPORT, REQUEST ASSISTANCE FROM RECEIVING AREA'S ADVANCED LIFE SUPPORT SERVICE.

STROKE / TIA or NEUROLOGICAL DEFICIT*





ADVANCED EMT ADVANCED EMT South COVID-19 ADDENDUM



PROTOCOL VARIANCES Guidance for COVID-19 Treatment and Transport



September 9, 2020

The following are updated variances for the 2020 SPEMS Protocols for dealing with COVID-19 in the pre-hospital setting and pertains to all levels of practice within the SPEMS Protocols.

These variances become effective immediately and will continue until the Medical Director deems that the COVID-19 threat has been eliminated.

******This variance replaces the variances dated March 23, 2020**

These protocol variances are:

- Insertion of the Suspected COVID-19 Treatment/Transport Algorithm dated 9/9/2020
- Insertion of the Non-Transport Guidance for Minor Cases of Suspected COVID-19 Algorithm dated 9/9/2020
- Insertion of the COVID-19 Treatment Supplements dated 9/9/2020
- To the equipment list of all levels of the SPEMS Protocols:
 - Add "2 Albuterol Metered Dose Inhalers (MDIs) (90mcg per activation) (As available)"
 - This medication should be added to each ambulance as soon as possible. This may be difficult to find but each EMS service should make every effort to find and stock this as soon as possible. It is imperative that patients with suspected COVID-19 NOT be given medications via a nebulizer.
 - It will not be a violation of protocol if an EMS service cannot find this medication. If unable to obtain, EMS services shall maintain documentation of backorder from their vendor.
 - Procedure should be followed as detailed in the COVID-19 Treatment Supplements dated 9/9/2020
 - Add "2 HEPA or Bacterial/Viral (B/F) Nebulizer Assemblies" (Optional)(As available)

Surgical Mask Recommendation: The American College of Emergency Physicians (ACEP) now recommends that EMS personnel wear surgical masks for all patient contact, if available. These masks can be used for an entire shift unless contaminated. All personnel should use an N-95 mask for all exposure to possible COVID-19 patients.

Again, these variances become effective immediately and will remain in effect until further notified by Medical Direction.

A copy of this page, both algorithms dated 09/09/2020 and the COVID-19 Treatment Supplements dated 09/09/2020 must be inserted into each official copy of the SPEMS Protocols.

Dr. Charles Addington II, D.O SPEMS Medical Director



If patient is transported to a facility:

as normal with one difference. Provide a covert war

- Radio Report: Proceed with report as normal with one difference. Provide a covert warning to the receiving facility of the high suspicion of CoVID-19. If cellular communication is not available then the driver of the unit should enter facility to advise of the situation while the patient and provider remain in the unit awaiting further instruction.
- At Destination: Once the patient has been moved over to facility bed and patient care has been transferred remove PPE and follow your agencies Standard Operating Procedures (SOPs) for disposal or reuse.
- Ambulance Disinfection: At a minimum, and while utilizing appropriate PPE, carefully clean/disinfect any surface contacted by the patient or provider before returning to service. Follow your agencies SOPs for equipment and ambulance disinfection.



Suspected CoVID-19 Treatment Procedure Supplements



- If available, the Metered Dose Inhaler (MDI) should be utilized with the above pictured closed system inhaler mask. This mask is used to help prevent the transmission of the virus by utilizing a closed system to limit aerosolization. This inhaler mask is assembled using a NRB or nebulizer mask, nebulizer spacer tube, two (2) ECG electrodes, and the MDI. Connect the corragated spacer tube to the mask and MDI and cover the two (2) exhaust ports on the mask with the ECG electrodes. (*Basic, Advanced, and Paramedic*)
- Instructional video: https://www.youtube.com/watch?v=8qaIk59u4mc&feature=youtu.be&app=desktop
- If MDIs are not available, **DO NOT** administer any nebulized bronchodilators unless done so utilizing a filtered nebulizer system (see following page) (*Basic, Advanced, and Paramedic*). Instead focus on basic airway management and the use of a BVM (*All levels*).
- CoVID-19 is considered as a droplet-precaution viral disease. However, droplets may be aerosolized by coughing, sneezing, or nebulized medication use (home nebulizer) and remain in the air for several hours. Use an N95 mask on yourself when making patient contact. If the patient is transported, apply a surgical mask to the patient to protect others. Do NOT use an N95 mask on these patients.
- If respiratory status continues to decline and the airway must be secured, **INTUBATION VIA THE DIRECT LARYNGOSCOPY TECHNIQUE IS NOT RECOMMENDED**. Instead, focus on BLS airway management to reduce the risk of personal contamination from respiratory droplets. Endotracheal intubation via indirect laryngoscopy (King Vision, Airtraq, etc...) or placement of a King Airway is an option, but only indicated for rare cases where BVM ventilation is inadequate.

"# †@) "u "h "o





Filtered Nebulizer Mask Assembly

0



- Traditional nebulizers are reported to increase droplet dispersion, causing uncertainty in use for COVID patients. A national shortage of Metered Dose Inhalers (MDI's) has led to difficulty obtaining them. However, administering a nebulized bronchodilator while utilizing a leak free filtered nebulized system has proven to be both effective and safer for healthcare professionals (*Basic, Advanced, and Paramedic*).
- If a nebulized bronchodilator is administered it **MUST** be done through a filtered nebulizer system that creates a seal around the patient's face throughout the entire treatment (if a mouth pieces is used in the place of a vent free mask the patient must maintain a seal around mouth piece throughout the entire treatment). It is recommended to use a vent free mask either strapped to the patient's face or held firmly in place by the EMS provider (*Basic, Advanced, and Paramedic*).
- There are different ways to create a leak free filtered nebulizer system. The above pictured technique is merely a suggestion. To make a safe and effective filtered nebulizer system it must be free of leaks where the system meets the patient's face or mouth. It must also utilize a HEPA (preferred) or a Bacterial / Viral (B/F) Filter located at the exhaust end of the system. If constructed adequately there should not be signs of the nebulized bronchodilator (fog) exiting anywhere from the filtered nebulizer system.
- Instructional video:
 k o uOnj U \)@U° j
- If MDIs or an effective filtered nebulizer system are not available, **DO NOT** administer any nebulized bronchodilators. Instead focus on basic airway management and the use of a BVM (*All levels*).
- If respiratory status continues to decline and the airway must be secured, **INTUBATION VIA THE DIRECT LARYNGOSCOPY TECHNIQUE IS NOT RECOMMENDED**. Instead, focus on BLS airway management to reduce the risk of personal contamination from respiratory droplets. Endotracheal intubation via indirect laryngoscopy (King Vision, Airtraq, etc...) or placement of a King Airway is an option, but only indicated for rare cases where BVM ventilation is inadequate.
- CoVID-19 is considered as a droplet-precaution viral disease. However, droplets may be aerosolized by coughing, sneezing, or nebulized medication use (home nebulizer) and remain in the air for several hours. Use an N95 mask on yourself when making patient contact. If the patient is transported, apply a surgical mask to the patient to protect others. Do NOT use an N95 mask on these patients.